

RESEARCH ARTICLE

Clinical Outcomes of Surgical Regenerative Treatment in Combination with Magnetic Laser Supportive Therapy of Peri-Implantitis

Hakobyan G^{1*}, Yessayan L², Khachatrayn G³ and Seyranyan A⁴

¹Professor, Head of Department of Oral and Maxillofacial Surgery Yerevan State Medical University, Armenia

²Professor, Head of Department of Therapeutic Dentistry, Yerevan State Medical University, Armenia

³Head of the Department of Dental Professional and Continuing Education Yerevan State Medical University, Armenia

⁴Assistant professor, Department of Oral and Maxillofacial Surgery Yerevan State Medical University, Dental Resident at Yerevan State Medical University, Armenia

*Corresponding author: Hakobyan G, Professor, Head of Department of Oral and Maxillofacial Surgery Yerevan State Medical University, Armenia, Tel: (+37410)271146, E-mail: prom_hg@yahoo.com

Citation: Hakobyan G, Yessayan L, Khachatrayn G, Seyranyan A (2020) Clinical Outcomes of Surgical Regenerative Treatment in Combination with Magnetic Laser Supportive Therapy of Peri-Implantitis. J Oral Infect Pathol 1: 103

Abstract

Objective: The objective this study is to evaluate the potential of magnetic-laser therapy as a supportive treatment of peri-implantitis.

Materials and Methods: The 34 patients suffering from Peri-Implantitis were selected for this study. The patients randomly divided into two groups;

18 patients first group received surgical conventional treatment,

16 patients second group received surgical conventional treatment and magnetic-laser application.

A total of 46 implants were treated with moderate peri-implantitis. Diagnostic parameters used to evaluate peri-implantitis include clinical indicators, Probing Pocket Depth (PPD), Bleeding On Probing (BOP), Marginal Bone Level (MBL) suppuration, mobility. Clinical and radiographical parameters were recorded before treatment (baseline) and at 3, 6 and 12, 36 months after therapy.

Results: Reduction PPD and BOP was observed in comparison with basic clinical measurements. The mean BOP in 34 patients before treatment of peri-implantitis was 2.5 ± 0.31 , after treatment, the first group of patients had mean 0.6 ± 0.1 , the second group had mean 0.4 ± 0.12 . The mean PPD in patients before treatment of peri-implantitis was 5.2 ± 0.24 , after treatment, the first group of patients had mean 3.9 ± 0.28 , the second group had mean 3.2 ± 0.17 .

The mean MBL concomitant bone level gain averaged was 1, 54 mm in first group and 2.35 mm in second group.

Stable clinical measurements PPD and BOP were demonstrated during the following 1,3 years.

Conclusion: Surgical regenerative treatment combined with magnetic-laser supportive therapy reliable method for treatment peri-implantitis and may be considered an adjunct to the conventional surgical treatments of peri-implantitis.

Keywords: Peri-Implantitis; Dental Implant; Regenerative Therapy; Magnetic-Laser Therapy

Abbreviations: PPD: Probing Pocket Depth; BOP: Bleeding On Probing; MBL: Marginal Bone Level; Er-YAG: Erbium-Doped Yttrium Aluminium Garnet

Introduction

Dental implantation is a successful method for treating partial or completely edentulous patients. However, according to various authors, in patients with implants, peri-implantitis is one of the late complications [1-4]. Peri-implantitis is a inflammatory disease of the tissues surrounding osseointegrated dental implants with varying degrees of peri-implant bone loss, increased pocket formation, purulence and eventually implant loss [5]. A consensus report identified the prevalence (5–10 year period) of peri-implantitis to be 28% to 56% of patients and 12% to 40% of implants [5,6]. Inflammatory process surrounding dental implants is represented in two common forms: peri-mucositis and peri-implantitis. The American Academy of Periodontology defines peri-mucositis is a reversible inflammatory response limited to the soft tissues surrounding an active oral implant. Peri-implantitis is a inflammatory response that involves loss of marginal bone around a functioning around osseointegrated implants, leading to the formation of a peri-implant pocket and loss of supporting bone [7].

The etiology of peri-implant diseases is characterized by various factors [8]. Risk factors include poor oral hygiene, no regular maintenance care after implant therapy, the volume of attached gingiva surrounding implant, the volume and quality of alveolar bone tissues, previous history of periodontal disease, para functional habits, smoking [9-16]. Bacterial biofilm, microbial colonization plays an important role in the etiology and development of peri-implantitis. Commonly found microorganisms are Gram-negative anaerobes, *Prevotella intermedia*, *Porphyromonas gingivalis*, *Aggregatibacter actinomycetemcomitans*, *Bacterioides forsythus*, *Treponema denticola*, *Prevotella nigrescens*, *Peptostreptococcus micros*, *Fusobacterium nucleatum* [17,18]. Excessive biomechanical forces may lead to high stress in the coronal bone-to-implant contact and thus lead to loss of osseointegration around the neck of the implant [19].

The impact of keratinized gingiva around dental implants has been controversially discussed [20]. In the development of peri-implantitis, the cement remaining after fixation of the restoration can also play a significant role [21]. Factors associated with the patient include systemic diseases, for example, diabetes mellitus, osteoporosis, prolonged treatment of corticosteroids, chemotherapy [22-23]. Forum and Rosen (2012) proposed classification for peri-implantitis depending on bleeding on probing, pocket depth and bone loss, peri-implantitis is divided into three categories: early, moderate, severe [24]. Strategies for the prevention and treatment of peri-implant diseases should be integrated into modern rehabilitation concepts in the field of oral implantology.

Peri-mucositis can be effectively treated with conservative methods. Prevention and treatment of peri-implantitis is extremely important because they lead to disintegration and loss of implants. For the treatment peri-implantitis, the etiological factor must be removed. Treatment of peri-implantitis is aimed at combating infection, decontamination of the surfaces implants, regeneration of lost tissue [25,26]. Various treatment methods are suggested in the treatment of peri-implantitis, non-surgical machining debridemen, chemotherapeutic disinfection, use of antibacterial agents, resective and regenerative surgical procedures, laser therapy, the employment of combination of lasers and surgical treatment [27-30]. However, there is no standard approach for the treatment of peri-implantitis, since with any of the treatment options complete elimination of inflammation is not achieved.

Nonsurgical therapy include with or without adjunctive local-release and systemic antibiotics [31-34]. In modern literature and expert assessments, mechanical processing of the implant is recommended, followed by early assessment and surgical intervention, decontamination of the implants surfaces using a variety of mechanical and chemical methods, then adjunctive systematic antibiotics [35]. Surgical treatments include a number of approaches (resective or regenerative) techniques, such as open flap debridement, augmentation and regenerative treatment. Surgical resection therapy for peri-implantitis is the recommended treatment option, however, due to the increase in postoperative recessions, this procedure is not recommended in aesthetic areas. Regenerative approaches include filling of the intraosseous peri-implant defect with a bone graft materials with a resorbable membranes and led to the most promising results [36,37].

Laser therapy is a modern therapeutic technique that can be effectively used as a complement to traditional mechanical therapy for the treatment of peri-implantitis. Diode lasers have been shown to have potent bactericidal effects based on this diode laser, carbon dioxide (CO₂) and Erbium Yttrium, Aluminum, Garnet (Er: YAG) lasers are suitable for irradiating [38-40]. Diode lasers have been shown to photobiomodulatory effects promoting wound healing and tissue regeneration. Diode lasers stimulate fibroblasts and osteoblasts, cause increased production of RNA messengers, which leads to significant collagen production during tissue healing [41-43]. The effect of a magnetic field on the surgical field also has an anti-inflammatory, analgesic, regenerating and accelerating tissue healing effect [44,45]. Such a similar therapeutic effect of laser radiation and magnetic field implies an increase in efficiency when used together.

The insufficient effectiveness of the proposed methods of treatment of peri-implantitis requires the improvement of surgical techniques, as well use innovative technology for the treatment of peri-implantitis. Based on this, research is needed to illustrate the clinical benefits of peri-implantitis treatment using magnetic-laser therapy. The objective this study is to evaluate the potential of magnetic-lazer therapy as a supportive treatment of peri-implantitis.

Materials and Methods

Thirty-four systemically healthy patients (16 females, 18 males, at a mean age 47,2years) suffering from peri-implantitis having pocket depth over 5mm in implants and having no mobility were selected for the study. Nine patients (12 implants) had a history of periodontal disease. All patients were informed of the study and received an informed consent form. The patients randomly divided into two groups;

- 18 patients first group received surgical conventional treatment,
- 16 patients second group received surgical conventional treatment and magnetic-lazer application.

46 implants were treated in total with moderate peri-implantitis. Clinical and radiological methods have been used to examine patients. Diagnostic parameters used to evaluate peri-implantitis include clinical indicators bleeding On Probing (BOP), Probing Pocket Depth (PPD), Marginal Bone Level (MBL) suppuration, mobility. Bleeding On Probing (BOP) is assessed as present if bleeding was evident within 30 seconds after the study or was not present if bleeding was not observed within 30 seconds after the study.

BOP indices were evaluated by the following criteria
0-no bleeding,
1-bleeding occurs no earlier than 30 seconds,
2-bleeding occurs in less than 30 seconds,
3-bleeding occurs when eating or brushing your teeth.

The degree of bleeding was assessed by the criteria
0.1-1.0 - mild inflammation,
1.1-2.0 - medium inflammation,
2.1-3.0 - severe inflammation.

The Probing Pocket Depth (PPD) was measured with a full millimeter with a manual periodontal probe from the edge of the mucosa to the bottom of the examined pocket. Indications for Marginal Bone Level (MBL) were evaluated by periapical radiographs (taken at the baseline diagnostic appointment). Before treatment (baseline) and at 3, 6 and 12, 24, 36 months after therapy clinical and radiographical parameters were recorded. Reduction BOP, PPD and MBL was observed in comparison with basic clinical measurements.

Treatment Protocols

The occlusion of all implant supported dental prosthesis was monitored and, if present, the extreme contacts were removed. Professional hygiene was carried out 7 days before the treatment, the patients were rinsed twice a day for 1 min with chlorhexidine 0.12%. The day before surgery, orally with a duration of 7-10 days, patients were prescribed systemic antibiotics (amoxicillin 500 mg and metronidazole 200 mg or augmentin 875 mg or ciprofloxacin 250 mg). For the selection of the most effective antibiotic for each case, microbial testing was performed.

Local anesthesia was accomplished by articain 4%. After local anesthesia, the superstructure was removed, incision was made around the neck of the implants and the flap of full thickness was raised to provide access to the defect of the peri-implant and the open surface of the implant. The abutment was removed and cover plugs were inserted in the implant. With titanium instruments granulation tissue was carefully removed in the bone defect. The implant surface is decontaminated with Air-Flow Perio Soft, successive topical applications of citric acid, 0.12% chlorhexidine, sterile physiological saline. Patients of the second group underwent magnetic-laser irradiation with a wavelength of 810 nm and a density of 100 mW for 30 seconds. Bone loss was evaluated intrasurgically, Bio-Oss was mixed with Gengigel hyaluronic acid preparation outside the mouth, and the peri-implant defect was filled. A bioresorbable collagen membrane Bio-Gide was placed over the filled defect. After bone grafting flaps were repositioned and sutured, wound healing was performed in a submerged. Patients were instructed to rinse twice a day for 1 minute for 2-3 weeks with chlorhexidine 0.12%.

After surgery the patients of the second group received magnetic laser irradiation 7 days with a wavelength of 810 nm and a density of 100 mW during 3 min. Healing periods occurred without complications, and with minimal postoperative discomfort. The sutures were removed 7-10 days after the surgery. To monitor healing, patients were observed for the first 4 weeks, and then at a three-month interval. Cover plugs of the implants were replaced with prosthetic abutments after 3 months of submerged healing and prosthetic components were installed after 1 week of soft tissue healing. Professional hygiene was conducted every six months. Effectiveness treatment was evaluated by the following criteria: (1) the absence of progressive loss of bone mass, (2) the absence of suppuration, (3) bleeding when probing for $\leq 50\%$ of sites and (4) Probing pocket depth $< 5\text{mm}$.

Statistical Analyses

Statistical analyses were performed using SPSS software ver. 22.0 (IBM, Armonk, NY, USA), and MedCalc program for Windows. To test the significance of variations in the BOP, PPD, MBL, the t-test was used. The minimum level of statistical significance was set at a value of less than 0.05.

Results

The diagnostic parameters of the two groups were comparable at baseline and after treatment. Radiologically increased or stable levels of the marginal bone compared with the baseline periapical x-rays is considered to be a treatment success. The mean initial both BOP and PPD of the groups was not significantly different ($p > 0.05$). Clinical evaluation of the results of treatment after 3, 6, 12 months showed reduction in both BOP and PPD were as compared with the baseline clinical measurements, more pronounced in the surgical treatment and magnetic-laser application method of treatment. The mean BOP in thirty-four patients before treatment of peri-implantitis was 2.5 ± 0.31 , after 6 months treatment. After 6 month treatment no statistically significant finding was observed in the mean BOP of both groups ($p > 0.05$), the first group of patients had mean 0.6 ± 0.1 , the second group had mean 0.4 ± 0.12 .

The mean PPD in thirty-four patients before treatment of peri-implantitis was 5.2 ± 0.24 . After 6 month treatment the volume of pocket depth in the first group (mean 3.9 ± 0.28) was significantly higher than the second group mean (3.2 ± 0.1) (Figure 1).




Figure 1: The mean BOP and PPD patients before and after 6 months treatment of peri-implantitis

The mean MBL concomitant bone level gain averaged was mean 1.74 mm in first group and mean 2.35 mm in second group 6 months after treatment. When considering the treatment benefit of both group, in patients received surgical conventional treatment and magnetic-lazer application, achieved better outcomes in terms of PPD reduction and MBL. At 1 year after initial treatment, stable clinical measurements of PPD and BOP were demonstrated and remained stable during the following three year. For the long-term stability of the treatment results, it is necessary for the patient to maintain good oral hygiene.

Discussion

Dental implants successful treatments for partial or full edentulous patients. One of the most commonly diagnosed complications of dental implants is peri-implantitis [8]. The etiology of peri-implantitis is multifactorial and treatment requires an integrated approach. Peri-implant diseases have always been associated with the biofilm, since it is generally accepted that peri-implantitis has a bacteriological etiology, decontamination of the implant surface is very important and the difference methods of the implant surface decontamination were used (such as powder air flow, saline flushing, citric acid, laser, hydrogen peroxide). Moderate and severe peri-implantitis will require surgical consideration. There are various surgical treatments methods (resective or regenerative). Surgical resection is used for peri-implantitis located in non-aesthetic sites [46]. Surgical regenerative therapy was carried out, using: (1) autogenous bone grafts alone(2), autogenous bone grafts covered by membranes, (3) xenogen bone grafts covered by membranes, (4) xenogen bone grafts, platelet-rich plasma covered by membranes [47]. Adjunctive peri-implant therapies, such as antibiotics, antiseptics, have been proposed to improve the non-surgical treatment options of peri-implantitis, however nonsurgical therapy is ineffective at treating peri-implantitis.

Different treatments methods have been suggested, but little reliable evidence exists, suggest-ing which could be the most effective one for the long-term treatments methods. Currently there is no universal approach therapy peri-implantitis [48]. The need to determine a standard treatment regimen for peri-implantitis was emphasized in the consensus report of the 8th European Periodontology Workshop [49]. Laser therapy is widely used in Dentistry. Laser therapy effective in periodontal treatment and there is ongoing debate regarding the efficacy of lasers in periodontal disease [50-53]. The American Dental Association's (ADA) and American Academy of Periodontology's (AAP) recommendations regarding their use in treating periodontal disease [54]. Since the role of periodontal pathogens in the etiology of peri-implantitis is indisputable, this indicates that anti-infection therapy for peri-implantitis should be consistent with the treatment strategy for periodontitis [55].

With the advent of laser technology, there is a growing tendency to use in implant dentistry. Now lasers have myriad applications in implant dentistry, which includes non-surgical therapy and is recommended in addition to the available conventional methods for the treatment of peri-implantitis [56]. The non-surgical treatment of peri-implantitis using Er:YAG laser showed in reduction peri-implant probing pocket depth and bleeding on probing. However, single course of treatment with the Er:YAG laser may not be adequate for achieving a stable therapy of peri-implantitis and that additional subsequent osseous regenerative procedures, might be required [57]. *In vitro* studies have shown that laser irradiation increases the release of bFGF from gingival fibroblasts [58]. With clinical use, this leads to an improved and accelerated wound healing [59]. Diode laser is used in peri-implant treatment to access the photochemical and photosensitive effects of laser. The main role of the laser in the treatment of peri-implants is its bactericidal effect. Magnetic-laser therapy combines the therapeutic factors widely used in modern medicine: magnetic field, low laser radiation and infrared light radiation.

The therapeutic effect of magnetic-laser therapy is manifested by immunomodulatory, anti-inflammatory, regenerative and enjoys from high bactericidal and detoxication effects [60-63]. This study describes clinical results of a magnetic-lazer therapy as a supportive treatment of peri-implantitis. Clinical and radiological evaluations of the results of treatment after 6 months showed reduction in both PPD and BOP as was compared with the baseline clinical measurements in both groups, a significant reductions

was shown in the second group of patients. At 1 year after initial treatment, stable clinical measurements of PPD and BOP were demonstrated and remained stable during the following three year. Magnetic-laser therapy has shown promising therapeutic effect in treatment of peri-implantitis. To achieve an optimal clinical result, at least 7 sessions of magnetic-laser therapy are necessary.

The clinical significance objectives of a magnetic-laser therapy approach are:

The reduce microorganisms on the implant surface,

1. The decrease BOP,
2. The reduce PPD,
3. To enhance self-performed oral hygiene and peri-implant health.

Magnetic-laser therapy is not only useful because of its bactericidal effect, but can also accelerate the regeneration processes in the peri-implant area. After 6 months x-ray examination demonstrated newly formed hard tissue was observed filling the defects around the implants and is considered to be a treatment success. For the long-term success of peri-implant treatment, constant dynamic monitoring and regular professional hygiene were performed.

The protective effect and slow absorption of hyaluronic acid provide reliable and predictable regeneration of augmented [64,65]. This barrier function of hyaluronic acid is very important in the process of wound healing. Our results suggest that the preparation of hyaluronic acid Gengigel is used in combination with Bio-Oss represents a good adjunctive treatment to conventional therapy of peri-implantitis.

The use of magneto-laser therapy for stabilization and decontamination of the affected surface of the implant has demonstrated promising results treating peri-implantitis. This combination of surgical and therapeutic treatment aims at improvement of the quality of regenerated bone structures. The results of this study indicated that a surgical procedure based on pocket elimination, bone grafting with grafts materials and hyaluronic acid Gengigel, magnetic-laser therapy accelerates the regeneration processes in the peri-implant area and was an effective therapy for treatment of peri-implantitis. It was found in the study that the combined magnetic -laser supportive therapy for implant surface debridement in peri-implantitis therapy were better at decreasing the PPD and MBL. Magnetic-laser therapy led to positive effects on clinical and radiologic parameters over the long-term subsequent period of time.

The surgical protocol for the treatment of peri-implantitis described in this article has shown positive results, therefore it is recommended as a simple and effective method of therapy.

Conclusion

The prognosis of the affected implant will depend on the early detection and treatment of peri-implant peri-implantitis. Supportive care should be given every 6 months. Surgical regenerative treatment combined with magnetic-laser therapy reliable method for treatment peri-implantitis. Magnetic-laser supportive therapy may be considered an adjunct to the conventional surgical treatments of peri-implantitis. The long-term success of peri-implant treatment requires constant dynamic observation, regular professional hygiene.

References

1. Sequeira V, Todkar MM, Abhyankar V, Fernandes G (2018) Prevention of Peri-Implantitis: A Narrative Review. *J Dent Oral Care Med* 4: 106.
2. Schwarz F, Derks J, Monje A, Wang HL (2018) Peri-implantitis. *J Clin Periodontol* 45: 246-66.
3. Rajesh KS, Spoorthi KH, Shashikanth Hegde, Arun Kumar MS (2015) Peri-implantitis 360. *Int J Dent Med Res* 1: 213-9.
4. Nguyen-Hieu T, Borghetti A, Aboudharam G (2012) Peri-implantitis: from diagnosis to therapeutics. *J Investig Clin Dent* 3: 79-94.
5. American Academy of Periodontology Task Force on Peri-implantitis (2013) Peri-implant mucositis and peri-implantitis: a current understanding of their diagnoses and clinical implications. *J Periodontol* 84: 436-43.
6. Linde J, Meyle J (2008) Peri-implant diseases: consensus report of the sixth European workshop of Periodontology. *J Clin Periodontol* 35: 286-291.
7. G Caton J, Armitage G, Berglundh T, Chapple ILC, Jepsen S, et al. (2018) A new classification scheme for periodontal and peri-implant diseases and conditions - Introduction and key changes from the 1999 classification. *J Clin Periodontol* 45: 1-8.
8. Zitzmann NU, Berglundh T (2008) Definition and prevalence of periimplant diseases. *J Clin Periodontol* 35: 286-91.
9. Alsaadi G, Quirynen M, Komarek A, Van Steenberghe D (2008) Impact of local and systemic factors on the incidence of late oral implant loss. *Clin Oral Implants Res* 19: 670-6.
10. Subramani K, Jung RE, Molenberg A, Hammerle CH (2009) Biofilm on dental implants: a review of the literature. *Int J Oral Maxillofacial Implants* 24: 616-26.
11. Korsch M, Obst U, Walther W (2014) Cement-associated peri-implantitis: A retrospective clinical observational study of fixed implant-supported restorations using a methacrylate cement. *Clin Oral Implants Res* 25: 797-802.
12. Smeets R, Henningsen A, Jung O, Heiland M, Hammächer C, et al. (2014) Definition, etiology, prevention and treatment of peri-implantitis-a review. *Head Face Med* 3: 34.
13. Mombelli A, Muller N, Cionca N (2012) The epidemiology of peri-implantitis. *Clin Oral Implants Res* 23: 67-76.
14. Karoussis IK, Salvi GE, Heitz-Mayfield LJ, Bragger U, Hammerle CH, et al. (2003) Long-term implant prognosis in patients with and without a history of chronic periodontitis: a 10-year prospective cohort study of the ITI Dental Implant System. *Clin Oral Implants Res* 14: 329-39.

15. Rocuzzo M, Bonino F, Aglietta M, Dalmasso P (2012) Ten-year results of a three arms prospective cohort study on implants in periodontally compromised patients. Part 2: clinical results. *Clin Oral Implants Res* 23: 389-95.
16. Ueno D, Nagano T, Watanabe T, Shirakawa S, Yashima A, et al. (2016) Effect of the keratinized mucosa width on the health status of periimplant and contralateral periodontal tissues: a cross-sectional study. *Implant Dent* 25: 796-801.
17. Heydenrijk K, Meijer HJ, Van Der Reijden WA, Raghoobar GM, Vissink A, et al. (2002) Microbiota around root-formed endosseous implants: A review of the literature. *Int J Oral Maxillofac Implants* 17: 829-38.
18. Ata-Ali J, Candel-Marti ME, Flichy-Fernandez AJ, Penarrocha-Oltra D, BalaguerMartinez JF, et al. (2011) Peri-implantitis: associated microbiota and treatment. *Med Oral Patol Oral Cir Bucal* 16: 937-43.
19. Naert I, Duyck J, Vandamme K (2012) Occlusal overload and bone/implant loss. *Clin Oral Implants Res* 23: 95-107.
20. Brito C, Tenenbaum HC, Wong BK, Schmitt C, Nogueira-Filho G (2014) Is keratinized mucosa indispensable to maintain peri-implant health? A systematic review of the literature. *J Biomed Mater Res B Appl Biomater* 102: 643-50.
21. Pette GA, Ganeles J, Norkin FJ (2013) Radiographic appearance of commonly used cements in implant dentistry. *Int J Periodontics Restorative Dent* 33: 61-8.
22. Javed F, Romanos GE (2009) Impact of diabetes mellitus and glycemic control on the osseointegration of dental implants: a systematic literature review. *J Periodontol* 80: 1719-30.
23. Chen H, Liu N, Xu X, Qu X, Lu E (2013) Smoking, radiotherapy, diabetes and osteoporosis as risk factors for dental implant failure: a meta-analysis. *PLoS One* 8: e71955.
24. Froum SJ, Rosen PS (2012) A proposed classification for peri-implantitis. *Int J Periodont Restor Dent* 32: 533-40.
25. Esposito M, Grusovin MG, Worthington HV (2012) Treatment of peri-implantitis: what interventions are effective? A Cochrane systematic review. *Europ J Oral Implantology* 5: 21-41.
26. Schwarz F, Hegewald A, John G, Sahn N, Becker J (2013) Four-year follow-up of combined surgical therapy of advanced peri-implantitis evaluating two methods of surface decontamination. *J Clin Periodontol* 40: 962-7.
27. Kotsovilis S, Karoussis IK, Trianti M, Fourmouis I (2008) Therapy of peri-implantitis: a systematic review. *J Clin Periodontology* 35: 621-9.
28. Roncati M, Lucchese A, Carinci F (2013) Non-surgical treatment of peri-implantitis with the adjunctive use of an 810-nm diode laser. *J Indian Soc Periodontology* 17: 812-5.
29. Mettraux GR, Sculean A, Bürgin WB, Salvi GE (2016) Two-year clinical outcomes following non-surgical mechanical therapy of peri-implantitis with adjunctive diode laser application. *Clin Oral Implants Res* 27: 845-9.
30. Okayasu K, Wang HL (2012) Decision tree for the management of peri-implant diseases. *Implant Dent* 20: 256-61.
31. Persson GR, Salvi GE, Heitz-Mayfield LJ, Lang NP (2006) Antimicrobial therapy using a local drug delivery system (Arestin) in the treatment of peri-implantitis: microbiological outcomes. *Clin Oral Implants Res* 17: 386-93.
32. Tang Z, Cao C, Sha Y, Lin Y, Wang X (2002) Effects of non-surgical treatment modalities on peri-implantitis. *Chin J Stomatology* 37: 173-5.
33. Salvi GE, Persson GR, Heitz-Mayfield LJA, Frei M, Lang NP (2007) Adjunctive local antibiotic therapy in the treatment of peri-implantitis II: Clinical and radiographic outcomes. *Clin Oral Implants Res* 18: 281-5.
34. Buchter A, Meyer U, Kruse-Losler B, Joos U, Kleinheinz J (2004) Sustained release of doxycycline for the treatment of peri-implantitis: Randomised controlled trial. *Br J Oral Maxillofac Surg* 42: 439-44.
35. Renvert S, Roos-Jansaker AM, Claffey N (2008) Non-surgical treatment of peri-implant mucositis and peri-implantitis: a literature review. *J Clin Periodontol* 35: 305-15.
36. Romeo E, Lops D, Chiapasco M, Ghisolfi M, Vogel G (2007) Therapy of peri-implantitis with resective surgery. A 3-year clinical trial on rough screw-shaped oral implants. Part II: Radiographic outcome. *Clin Oral Implants Res* 18: 179-87.
37. Berglundh T, Wennström JL, Lindhe J (2018) Long-term outcome of surgical treatment of peri-implantitis: A 2-11-year retrospective study. *Clin Oral Implants Res* 29: 404-10.
38. Trivedi AR, Patel VG, Gupta SS, Jathal BS (2019) Treatment of Peri-implantitis with Implantoplasty and Diode Laser. *Int J Clin Med Imaging* 5: 49-53.
39. Guo-Hao Lin, Fernando Suárez López del Amo (2018) Hom-Lay Wang Laser therapy for treatment of peri-implant mucositis and peri-implantitis: An American Academy of Periodontology best evidence review. *J Periodontol* 89: 766-82.
40. Behdin S, Monje A, Lin GH, Edwards B, Othman A, et al. (2015) Effectiveness of laser application for periodontal surgical therapy: systematic review and meta-analysis. *J Periodontol* 86: 1352-63.
41. Lucas Freitas de Freitas, Michael R Hamblin (2016) Proposed Mechanisms of Photobiomodulation or Low-Level Light Therapy. *IEEE J Sel Top Quantum Electron* 22: 7000417.
42. de Abreu Chaves ME, de Araujo AR, Cruz Piancastelli AC, Marcos Pinotti M (2014) Effects of low-power light therapy on wound healing: LASER x LED*. *An Bras Dermatol* 89: 616-23.
43. Vartika Kathuria, Jatinder Kaur Dhillon, Gauri Kalra (2015) Low Level Laser Therapy: A Panacea for oral maladies. *Laser Ther* 24: 215-23.
44. K Marycz, K Kornicka, M Röcken (2018) Static magnetic field (SMF) as a regulator of stem cell fate-new perspectives in regenerative medicine arising from an underestimated tool. *Stem Cell Reviews* 14: 785-92.
45. Darendeliler MA, Darendeliler A, Sinclair PM (1997) Effects of static magnetic and pulsed electromagnetic fields on bone healing. *Int J Adult Orthodontics and Orthognathic Surg* 12: 43-53.
46. Chen S Darby I (2003) Dental implants: maintenance, care and treatment of peri-implant infection. *Aust Dent J* 48: 212-20.
47. Samizade S, Kazemian M, Ghorbanzadeh S, Amini P (2015) Peri-implant diseases: Treatment and management. *Int J Contemp Dent Med Rev* 2015: 6.
48. Schou S, Berglundh T, Lang NP (2004) Surgical treatment of peri-implantitis. *Int J Oral Maxillofac Implants* 19: 140-9.
49. Berglundh T, Stavropoulos A, Working Group 1 of the VIII European Workshop on Periodontology (2012) Preclinical in vivo research in implant dentistry. Consensus of the eighth European workshop on periodontology. *J Clin Periodontol* 39: 1-5.
50. Jeffrey D Pope, Jeffrey A Rossmann (2015) Laser Therapy in Periodontal Treatment. *Decisions in Dent*.
51. C Ren, C McGrath, L Jin, C Zhang, Y Yang (2017) The effectiveness of low-level laser therapy as an adjunct to non-surgical periodontal treatment: a meta-analysis. *J Periodontol Res* 52: 8-20.

52. Gregg RH, Mc Carthy D (2002) Laser periodontal therapy for bone regeneration. *Dent Today* 21: 54-59.
53. American Dental Association (ADA) (2015) Council on Scientific Affairs: Statement on Lasers in Dentistry.
54. Costa FO, Ferreira SD, Cortelli JR, Esteves Lima RP, Cortelli SC, et al. (2019) Microbiological profile associated with peri-implant diseases in individuals with and without preventive maintenance therapy: a 5-year follow-up. *Clin Oral Invest* 23: 3161-71.
55. Lin GH, Suarez Lopez Del Amo F, Wang HL (2018) Laser therapy for treatment of peri-implant mucositis and peri-implantitis: An American Academy of Periodontology best evidence review. *J Periodontol* 89: 766-82.
56. Schwarz F, Bieling K, Bonsmann M, Latz T, Becker J (2006) Nonsurgical treatment of moderate and advanced periimplantitis lesions: a controlled clinical study. *Clin Oral Investig* 10: 279-88.
57. Almeida-Lopes L, Rigau J, Zangaro RA, Guidugli-Neto J, Jaeger MM (2001) Comparison of the low level laser therapy effects on cultured human gingival fibroblasts proliferation using different irradiance and same fluence. *Lasers Surg Med* 29: 179-84.
58. Schwarz F, Aoki A, Sculean A, Becker J (2009) The impact of laser application on periodontal and peri-implant wound healing. *Periodontol* 2000 51: 79-108.
59. Fouda A, Refai H, Mohammed N (2013) Low level laser therapy versus pulsed electromagnetic field for inactivation of myofascial trigger points. *Am J Res Commun* 1: 68-78.
60. Kathuria V, Dhillon JK, Kalra G (2015) Low Level Laser Therapy: A Panacea for oral maladies. *Laser Ther* 24: 215-23.
61. Sculean A, Schwarz F, Becker J (2005) Anti-infective therapy with an Er: YAG laser: Influence on peri-implant healing. *Expert Rev Med Devices* 2: 267-76.
62. Ashnagar S, Nowzari H, Nokhbatolfoghahaei H, Zadeh BY, Chiniforush N (2014) Laser Treatment of Peri-Implantitis: A Literature Review. *J Lasers Med Sci* 5: 153-62.
63. Parveen Dahiya, Reet Kamal (2013) Hyaluronic Acid: A Boon in Periodontal Therapy. *N Am J Med Sci* 5: 309-15.
64. Pagnacco A, Vangelisti R, Erra C, Poma A (1997) Double-blind clinical trial versus placebo of a new sodium-hyaluronate-based gingival gel. *Attual Ter In* 15: 1-7.
65. Ballini A, Cantore S, Capodiferro S, Grassi FR (2009) Esterified hyaluronic acid and autologous bone in the surgical correction of the infra-bone defects. *Int J Med Sci* 6: 65-71.