

RESEARCH ARTICLE

Isoetes Pantii Complex is Evolving with New Basic Chromosome Number

Goswami HK¹ and Majumdar J^{2*}¹Ph.D. Retired and Visiting Professor of Botany and Genetics, 24- Kaushalnagar, Misrod, Bhopal (MP), India²Department of Biological Sciences, Burdwan Town School, Burdwan, West Bengal, India

*Corresponding author: Mazumdar J, Department of Biological Sciences, Burdwan Town School, Burdwan-713101, India, E-mail: jaideepmazumdar10@gmail.com/hitkishoregoswami@yahoo.in

Citation: Goswami HK, Mazumdar J (2019) *Isoetes Pantii* Complex is Evolving with New Basic Chromosome Number. J Adv Plant Sci 2: 107

Abstract

A new species, *Isoetes pantii* HK Goswami and BS Arya was described more than fifty years ago from Narsingharh in Central India. Many of the rare features of this species, like heterosporous microsporangia (now known as heterosporangia), three types of microspores (alete, monolete, trilete) and several plants possessing $2n=48$ chromosomes (basic $n=12$) have been repeatedly confirmed since then. Continued search for *Isoetes* plants from several localities further yielded two new species. Plants of both these species resembled *I. pantii* in having round tuberculation megaspores but exhibited $n=12$ series and showed $2n=60$ chromosomes in roots. However both these species have exhibited extremely rare features. *I. fuchsii* sp Nov possessed 25% megaspores with spines on the proximal face in only one pyramidal area, a feature never observed so far. Plants of another species published earlier as *I. pantii* var *hybrida* (hereunder being revised and named as *Isoetes narsingharhensis* sp Nov) were very small (less than 10cms) and exhibited a very strange mixed/hybrid trait on exine ornamentation of megaspores; megaspores possess both reticulate exine as well as clearly demarcated round tubercles. Such mixed exine ornamentation has never been described in the botanical literature. Since both these species were not properly published we wish to describe them again and confirm the validity of *I. fuchsii* and also propose a new name *Isoetes narsingharhensis* sp Nov for *I. pantii* var *hybrida*. We also assert that the *Isoetes pantii* complex is essentially characterized by round tuberculation megaspores but with chromosome counts based on $n=12$ chromosome series. As far as known, no species of *Isoetes* in the world flora possess multiples of $n=12$ chromosomes; infact all other species possess $n=11$ chromosome series. While all essential evidences already published, *Isoetes pantii* is a confirmed sexual species (male and female gametophytes and developing sporophytes have been well described earlier) we have so far not found any sexual stage either in *I. fuchsii* or in newly named *I. narsingharhensis* sp nov hence we presume these plants to be sterile hybrid segregates constituting *I pantii* complex.

Keywords: Cytology; India; Isoetaceae; *Isoetes pantii* Complex; New Species

Introduction

The genus *Isoetes* Linnaeus (1753: 1100; Linneus filius C.) [1,2] (Isoetaceae) is represented by about 250 species in the World (PPG I 2016). [3]. In a recent revision Fraser-Jenkins, *et al.* [4] published that only four species occur in India. Certainly such a statement is oversimplification of the fact and as such cannot be accepted. This treatment has not been accepted by many workers; however denial of genuine facts ignoring excellently published papers. In the meantime, Troia, *et al.* [5] observed that the taxonomy of *Isoetes* in India is unclear. The hard fact on the contrary is that there are many more species validly reported from India [6-10] established on the basis of spore morphology (as strongly confirmed by Turland, *et al.*, Art. 39.1; Turland, *et al.* Art. 40.1) and chromosome numbers [11]. The criticism of recognition of *Isoetes pantii* HK Goswami and BS Arya in particular was once advocated by those workers who do not give any cognizance to chromosome constitutional variation (position of centromere on the chromosomes) and chromosome number. *I. pantii* has been shown to possess sex chromosomal variation in meiosis and mitotic studies observed since early days of 1970 and excellent demonstration by shape, size, meiotic behaviour and molecular genetic studies have been well documented [7,12-17]. This has been proved by repeated studies since then that *I. pantii* has evolved with basic chromosomes $n=12$ and several variations have been described with stable count of $2n=48$ chromosomes among many plants [15,18,19]. Another relevant feature with great morphological impact worth emphasizing regarding *Isoetes pantii* Goswami & Arya (1970: 30) [20] (Figure 1A), has been the development of three kinds of microspores (alete, monolete and trilete) in the microsporangia along with highly unusual large megaspores inside within the same microsporangia [15,18-21]. These all features have been repeatedly observed since 1966 onwards and are excellent acquisition within the genome of *Isoetes pantii*. No species in

the world flora of the genus *Isoetes* has ever shown a combination of these traits which decidedly indicate strong genetic divergence from other species. Elsewhere, molecular genetic studies have been published to opine as to why very unusual megaspores resembling fossil lycopod megaspores develop inside the microsporangium of *I. pantii* and what mechanism might be responsible for triggering female “genes” inside the male sporangium (?). This switching on mechanism might be due to one of the putative parental genes (*I. sampathkumaranii* LN Rao) becoming active because megaspores inside the microsporangia possess reticulate exine while megaspores inside the mega sporangium are round tuberculation. These large megaspores are fertile and produce female gametophyte [15,18,22-26]. The presence of sex chromosomes reported in *I. pantii* [7,13,15,16,26-29] have been well cited thereby establishing an altogether different path of chromosomal evolution (Slides of chromosome preparations; spore morphologies and microtome sections along with plant materials were shared with Late Professors DD Pant, Fuchs-Eckert during their visits to Bhopal and excellent photographs have been published over these years). Follow-up cytogenetic studies on *Isoetes* populations have had further yielded two new species. Plants of both these species resemble *I. pantii* in having round tuberculation megaspores, possess (n=12 series) 2n=60 chromosomes in roots and have also exhibited extremely rare features. *I. fuchsii* sp Nov possessed 25% megaspores with spines on the proximal face a feature never observed so far. Plants of another species described earlier as *I. pantii* var *hybrida* (hereunder being revised and named as *Isoetes narsingharhensis* sp Nov) were very small (less than 10cms) and exhibited a very strange mixed/hybrid trait on exine ornamentation of megaspores; megaspores possess both reticulate exine as well as clearly demarcated round tubercles. Since both these species were not properly published earlier, we wish to describe them again and confirm the validity of *I. fuchsii* and also propose a new name *Isoetes narsingharhensis* sp Nov for *I. pantii* var *hybrida*. We also assert that the *Isoetes pantii* complex is essentially characterized by round tuberculation megaspores but with chromosome counts based on n=12 chromosome series. All other species of the genus possess n=11 chromosome series. While all essential evidences already published, *Isoetes pantii* is a confirmed sexual species (male and female gametophytes and developing sporophytes have been well described earlier) we have so far not found any sexual stage either in *I. fuchsii* or in newly named *I. narsingharhensis* sp Nov hence we presume these plants to be sterile hybrid segregates constituting *I. pantii* complex.

Materials and Methods

While the major localities and ponds for collections were at Narsingharh (MP) India, extensive collections (since 1966 onwards) and population based comparative morphological, anatomical, biochemical, chromosomal, and molecular genetic studies have been carried out (Table 1) (Figure 1, 2) from various parts of India and series of papers have been published (some of them among present references) dealing with many species Viz. *Isoetes coromandelina*, *I. pantii*, *I. indica*, *I. sampathkumaranii*, *I. dixitei*, *I. panchananii*, and a few more which are under repeated scrutiny [7,8]. All related methodologies have already been published. Briefly, plants were collected from respective localities and young roots were put in 8-hydroxyquinolene for four hours and then transferred to 45% acetic acid. After a few hours roots were squashed and stained with 1.5% acetic-orcein and immediately examined under a light microscope attached with camera. Good and well stained metaphases were photographed.

Features	<i>I. pantii</i>	<i>I. fuchsii</i>	<i>I.narsingharhensis</i>	<i>I.coromandelina</i>	<i>I.sampathkumaranii</i>
Rhizome	3 lobed, but one lobe always smaller than the other two	3-4 lobed	2 lobed	3-4 lobed	2 lobed
Velum	Young sporangia may have small velum, but absent on maturity	absent	covering ¼ of sporangium	absent	covering ½ of sporangium
Megaspore	trimorphic	trimorphic	dimorphic	dimorphic	dimorphic
Megaspore exine ornamentation	tuberculate	tuberculate	Mixed tuberculate,-- Reticulate (Figure 1F)	tuberculate	reticulate
Spine in megaspore	absent	Present on proximal face at one pyramidal area (Figure 1E)	Rarely, small spine present on abortive spores (Figure 1G)	absent	absent
Chromosomes	2n= 24, 36, 48 (including sex chromosomes and two B chromosomes)	2n=60 with 1 or 2 B chromosomes	2n=60 with 1 or 2 B chromosomes	2n=33, 44 + 1B	2n=66 + 1 B

Table 1: Comparative morphology of *I. pantii* complex with related species




Figure 1: (A) Habit of plants: *Isoetes fuchsii* (left), *I. narsinghgharensis* (middle), *I. pantii* (right); (B, C) Megaspores of *I. fuchsii* from mega sporangium stained with safranin showing joint spores with small spiny outgrowths all around periphery. Some spores also exhibit large spine emerging from the round tubercle; (D) A megaspore of *I. fuchsii* showing spine emerging from feebly developed tubercle in one pyramidal area; (E) Plants of *I. narsinghgharensis*; (F) Megaspores of *I. narsinghgharensis* from mega sporangium showing mixed occurrence of reticulate exine with sporadic round tubercles, a combination never so far known on any exine ornamentation; (G) An ill developed young megaspore of *I. narsinghgharensis* with prominent spine under SEM




Figure 2: Mitotic chromosomes of *Isoetes fuchsii* $2n=60$; the lower black arrow indicates an acrocentric chromosome in association with a chromosome (telocentric one) and the upper one solid black arrow is in the vicinity of a B chromosome (detailed significance with differential staining is still unpublished)

The inherent intention of this paper is to recognize the taxonomic status of *I. pantii* complex suspected earlier [12,13] and recognize proposed these two species, *Isoetes x fuchsii* I Bhu, HK Goswami, US Sharma and AK Bajpai, sp. Nov and *I. narsingharhensis* H K Goswami and J Mazumdar, sp Nov as the major constituent.

Observations and Comments

Taxonomic account

***Isoetes pantii*:** HK Goswami and BS Arya (1971: 30) (Figure 1A)

Type: INDIA. Madhya Pradesh: Rajgarh, Narsingharh, HK Goswami GA: IA (holotype BM!; isotypes DD!, Herbarium of Botany Department Allahabad University, Allahabad! Herbarium of Botany Department Government College, Narsingharh! Herbarium of Botany Department Govt. Science College, Gwalior!; plants and slides showing mixed occurrence of micro and megaspores have also been deposited at Kew and BME, England.

Description: Plants 24-60cm (Figure 1A); details of diagnosis of the species have already been described [7,14,19-21]. Pertinent points worth mentioning are that plants possess $2n=48$ chromosomes including sex chromosomes and 02 B (dot) chromosomes [7,12-14,18,19] are fertile and possess hybrid features of exine ornamentation on megaspores. Peculiarly, megaspores developing inside heterosporangia [21,26] show reticulate exine like megaspores of *I. sampathkumaranii* and megaspores in megasporangia are round tuberculate resembling *I. coromandelina*. Since both *I. coromandelina* and *I. sampathkumaranii* grow intermixed in the ponds at Narsingharh (MP, India) several morphological, biochemical and molecular genetic studies on genomic DNA procedures including Southern blot experiments on many related species were conducted. The hypothesis that both the above species should have crossed in nature [13,23,30,31] and the variant plants in ponds being observed since 1966 and thereafter, might be actually natural segregates of the natural hybridization. Furthermore, as germination of megaspores to produce female gametophytes and young embryos have already been published earlier (23, 24,25, 26) we strongly believe that *I. pantii* is a fertile hybrid.

Etymology: Named in honour of renowned Indian botanist Prof. Divya Darshan Pant (1919-2001).

Chromosome number: Root tip mitosis shows $2n=48$ including sex chromosomes and two B chromosomes; A few plants with $2n=36$ including B-chromosome (1-2). These plants also show typical features of *I. pantii*.

Ecology: Plants are submerged in water body.

Fertility: August - September–October.

Distribution: So far only known from India (Narsingharh and Rampura in Madhya Pradesh).

Isoetes x fuchsii

Isoetes x fuchsii I Bhu, HK Goswami, US Sharma and AK Bajpai, sp Nov (Figure 1A).

***Isoetes fuchsii*:** Bhu, *et al.* (2001: 12)[32] invalid publication on account of lack of Latin diagnosis, needed at that time.

Type: INDIA. Madhya Pradesh: Guna, Maksoodangarh, H.K. Goswami HKG-2000 (holotype Herbarium of Society of Bionaturalists, Bhopal!; Isotypes K!, Herbarium of Bioscience Department (pteridophyte section) South Gujrat University, Surat!).

Description: Plants measure 20-30cm (Figure 1A), rhizomorph 3 or 4 lobed; leaves measure 20-26cm, almost of equal in length; velum absent; megaspores with round tubercles; mega sporangia many, containing about 400 to 490 megaspores; megaspores trimorphic with round tubercles, large megaspores white when dry, ash coloured when wet, diameter range from 350-400 μ m, 374 μ m (average of 50 megaspores), globose to elongated in shape; tri-radiate ridge often straight, rarely sinuous; proximal face with 3-4 round tubercles, distal face many small size tubercles; medium size spores, tubercles measure 200 to 300 μ m in diameter; proximal face with 1-3 uneven tubercles, distal face possesses 20 or more tubercles (Figure 1B and C) about 20% spores show unusually significant spine emerging from the pyramidal area (Figure 1E) small megaspores not many in number even less than 5% among spores measure less than 100 μ m in diameter; proximal face with one large tubercle having a nipple like protuberance and the corresponding small spores may show depression/ pit in the middle of tubercle [32]; microsporangia not present in all plants; microspores dimorphic, only monolete and trilete spores have been observed, microspores measure 20-29 μ m with wide perispore; no megaspores have been observed in microsporangia. Root tip mitosis shows $2n=60$ chromosomes (Figure 2).

Etymology: This new species is named in honour of Swiss botanist Prof. Hans Peter Fuchs-Eckert (1928-1999) for his noteworthy contributions to the genus *Isoetes* and also approving *I. fuchsii* as new species.

Chromosome number: Root tip mitosis shows $2n=60$ chromosomes (Figure 2); meiotic studies did not reveal excellent preparations, but no laggards were observed at any stage.

Diagnosis: *I. fuchsii* possesses trimorphic megaspores with round tubercles and belong to Section Palustres (=Tuberculatae). About 15–20 % megaspores have large spines, a feature never reported so far in any species of the genus *Isoetes* in the world flora. Among Small megaspores, some have pits on the middle of round tubercle and some have nipple like protuberance as if the two had attached earlier during sporogenesis.

I. fuchsii possesses $2n=60$ chromosomes following the path of *I. pantii* series of $2n=24; 36, 48$; which are based on $n=12$ series of chromosome evolution.

Distribution: So far only know from India, in water bodies in Maksoodangarh, Guna and low ponds in forest area around Maksoodangarh.

Ecology: Plants are normally submerged or in marshy area toward margin of the water body.

Fertility: September- October –November.

Isoetes narsingharhensis

Isoetes narsingharhensis HK Goswami and Mazumdar sp Nov (Figure 1A,D,F and G)

***Isoetes pantii*:** Goswami and Arya (1971: 30) var *hybrida* Goswami (2004: 152) nom. inval.

Type: INDIA. Madhya Pradesh: Rajgarh, Narsingharh, H.K. Goswami HKG/ IS-2004 (holotype Herbarium of Society of Bionaturalists, Bhopal!; isotypes Herbarium of Bioscience Department (pteridophyte section), South Gujrat University, Surat!, Bapalal Vaidya Botanical Research Centre, Surat!; Isotypes K!).

Description: Plants attain height up to 10cm (Figure 1A, D), rhizome bi-lobed; rarely 3 lobes; sporangia do not show velum; mega sporangia oval to round containing 47 & 233 megaspores in outer and middle sporangia respectively; dimorphic, tri-radiate, large megaspores 100-250 μ m in diameter, proximal face possesses 3-8 small round tubercles; on distal face exine typically reticulate or wavy; few spores may possess spiny outgrowth (Figure 1G); small megaspores tri-radiate 50-95 μ m in diameter, both proximal face and distal face have small round tubercle like structures; each sporangium consists of 5 to 6% anucleate and or abortive deshaped spores; presence of both tubercles on proximal face and wavy, reticulate exine on the distal face of large spores along with presence of abortive spores as a regular feature (Figure 1F) decidedly indicates this to be a natural segregate; microsporangia have not been found so far.

Diagnosis: Unique combination of features: bi-lobed rhizome and tuberculation and reticulate megaspores in same mega sporangium (Figure 1F) distinguish it from other species of *I. pantii* complex.

Etymology: This species is named after the locality Narsingharh (Madhya Pradesh, India) where it was first discovered. Earlier epithet “*hybrida*” was derived from the belief that this new taxon was a natural hybrid between *I. coromandelina* and *I.sampathkumaranii* Rao (1944: 287)[33].

Distribution: So far only known from in India (Narsingharh and Bhopal in Madhya Pradesh).

Ecology: Plants grow as amphibious in periphery of water body.

Fertility: October –November- December

Discussion

Taxonomic relevance

Unfortunately *I. fuchsii* was published without a latin diagnosis [11] and *I. pantii* var *hybrida* without a holotype citation [11] and thus invalid as pointed out by Fraser-Jenkins, *et al.* [4]. These names are republished here for inclusion in the World flora of *Isoetes*, along with synopsis of *I. pantii* complex. So also the publication of *I. pantii* var. *Hybrida* (Goswami, 2004) [8] was pointed out to be invalid as per taxonomic practices. Hence we republish these two species along with a brief background of evolutionary relevance.

I x fuchsii Bhu, *et al.*

A new species named after Professor Fuchs-Eckert was published [32] which showed very peculiar features of possessing nearly 20-25% megaspores showing large spines and root tips showing $2n=60$ chromosomes, corresponding to the chromosomal path of evolution based on $n=12$ chromosomes. Since the latin diagnosis was not added, which was essential at that time of publication, the validity of species was obviously questioned. We therefore revise and offer this text so as to properly validate *I fuchsii* Bhu, *et al.* As already mentioned plants have been deposited at various herbaria. This species is a natural sterile hybrid

Isoetes x narsingharhensis Goswami and Mazumdar

Ponds in Narsingharh (*pachtalai* pond in particular) and in adjoining villages plant collections had initially identified three species; *I. coromandelina*, *I sampathkumaranii* and *I. pantii* as well as certain variants [12,18]. Our follow-up studies also recorded the growth of small plants confined to the periphery of the same pond at Narsingharh which were smaller but typically characterized by $2n=60$ chromosomes and very strangely megaspores exhibited both reticulate exine as well as presence of round tubercles. Such a feature has never been observed in any species so far. Assuming this to be a clear hybrid, this was published as *I. pantii* var.*hybrida* [8]. Now, on the basis of repeated observations and ascertaining about the permanency of all specific traits, we have renamed these plants as *I. narsingharhensis* (Figure 1D and F).

Evolutionary significance of “*pantii* complex”

Since August 1966 more than a dozen papers on morphological, anatomical, population cytogenetic screening and genomic assays have been published [6,11-13,19,20,24,27]. The discovery of *I. pantii* has led to establish a new series of chromosomal evolution [15-17,34] and three taxa with basic chromosome number $n=12$, have been observed to constitute the assemblage of three species *I. pantii* ($2n=48$), *I. fuchsii* ($2n=60$) and *I. narsinghgarhensis* with $2n=60$ chromosomes. Additionally, these species have genes for exine ornamentation from *I. coromandelina* as well as from *I. sampathkumaranii*; megaspores developing inside the microsporangia of *I. pantii* possess reticulate and webbed exine (like *sampathkumarinii*) while the megaspores inside the mega sporangia of the same plant possess round tubercles (like, *coromandelina*). This is relevant to record here that *I. narsinghgarhensis* megaspores possess mixed occurrence of reticulate exine and round tubercles on the distal as well as on proximal face; which is an extremely rare morphological combination of exine ornamentation.

Needless to mention, these species can not be considered to be cytotypes because each one has distinct features of taxonomic importance. These species appear to have evolved on a chromosomal path totally different and have assemblage of such DNA sequences which are endowed with rare sequences comparable to many other organisms (details are out of present context; see Goswami & Chandorkar, Goswami & Bhu, Bajpai & Goswami [22,24,31,26]. Obviously, then “*panti* complex” members have got to have chromosomes based on, $n = 12$ series.

Summary

Besides observing and ascertaining a new series of chromosomal evolution based on $n=12$ [15-17], we have also encountered evolution of new traits over these years. Prominent among these are, large spines on the proximal face of some megaspores (Figure 1) on one pyramidal area of *I. fuchsii* (Figure 1B and C) and mixed occurrence of reticulate exine as well as tubercles on the distal and proximal faces of *I. narsinghgarhensis* (Figure 1D and F).

Conservation status

Ever increasing human interference on lakes and ponds and invasion of weeds like *Ipomea*, *Lantana camara* etc on marshy area around water bodies are the commonest incidences noticed at all places thereby reducing the populations of these small quillworts [25]. Furthermore, villagers have started ploughing on these sites therefore survival of flora particularly growing on catchment marshy areas are getting terribly reduced. Until some institutional conservation strategies are not opted, it would be difficult to conserve these evolutionary important species [26]. HKG has been trying in vain to convince local villagers, local teachers and students of botany for protection of small lakes?

Acknowledgements

HKG is grateful to the Late Prof. DD Pant (Allahabad) and Late Mr. Clive Jermy (London), for examining slides, materials and helping with photographs during 1970-1980s and helping with references on *Isoetes pantii* and Late Prof. Fuchs Eckert for respective approvals and helpful suggestions. The plant material and slides of sporangia have also been personally submitted by HKG and we are grateful to Dr. Alison Paul for her help at BM in London and Sally Dawson at Royal Botanic Gardens at Kew.

References

1. Linnaeus C (1753) Species plantarum. L. Salvius, Stockholm.
2. Linnaeus filius C (1782) Supplementum plantarum. Impensis Orphanotropei, Brunsvigae.
3. PPG I (2016) A community-derived classification for extant lycophytes and ferns. J Syst Evol 54: 563-603.
4. Fraser-Jenkins CR, Gandhi KN, Kholia BS, Benniamin A (2016) An Annotated Checklist of Indian Pteridophytes Part-1 (Lycopodiaceae to Thelypteridaceae). Bishen Singh Mahendra Pal Singh.
5. Troia A, Pereira JB, Kim C, Taylor WC (2016) The genus *Isoetes* (Isoetaceae): a provisional checklist of the accepted and unresolved taxa. Phytotaxa 277: 101-45.
6. Fuchs-Eckert HP (1992) Supplementum and indicem isoëtaliū. Bionature 12: 99-159.
7. Goswami HK (1996) Three decades with *Isoetes*. Indian Fern J 19: 51-61.
8. Goswami HK (2004) Some observations on *Isoetes pantii* complex: New segregate in nature. Indian Fern J 21: 150-57.
9. Mazumdar J, Rajput K, Patil S (2017) (2565) Proposal to reject the name *Isoetes sahyadrii* (Isoetaceae). Taxon 66: 1470.
10. Singh SK, Shukla SK, Dubey NK, Shukla PK (2018) *Isoetes xgopalkrishnae* (Isoetaceae), a new interspecific sterile hybrid from central India. Phytotaxa 362: 68-76.
11. Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, et al. (2018) (edition) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. Regnum Vegetabile 159. Glashütten: Koeltz Botanical Books.
12. Goswami HK (1975a) The morphogenetics of mixed sporangia in *Isoetes*. J Indian Bot Soc 54: 210-8.
13. Goswami HK (1975b) Chromosome studies in natural populations of *Isoetes pantii* with heterosporous sporangia. Cytologia 40: 543-51.
14. Goswami HK (2011) Population cytogenetic studies confirm X-Y mechanism in *Isoetes pantii* (Isoetaceae: Pteridophyta). Bionature 31: 13-22.
15. Bhu I, Goswami HK (1990) A new line of chromosomal evolution in *Isoetes*. Bionature 10: 45-53.
16. Jermy AC (1991) Isoetaceae. In, Pteridophytes and Gymnosperms (Eds: Kramer KU & Green PS, Springer-Verlag, Berlin
17. Liu X, Gituru WR, Wang QF (2004) Distribution of basic diploid and polyploid Species of *Isoetes* In East Asia. J Biogeogr 31: 1239-50.

18. Goswami I, Goswami HK (1986) Genetics of natural variants VI. Some abnormal spores of *I. pantii* Goswami & Arya resemble fossil lycopods. *Bionature* 6: 23-28.
21. Goswami HK, Arya BS (1968) Heterosporous sporangia in *Isoetes*. *British Fern Gaz* 10: 39-40.
19. Goswami HK, Arya BS (2012) *Isoetes* × *pantii* (Isoetaceae: Pteridophyta): A Review. *Nelumbo* 54: 187-92.
20. Goswami HK, Arya BS (1970) A new species of *Isoetes* from Narsingharh, Madhya Pradesh. *J Indian Bot Soc* 49: 30-7.
22. Goswami HK, Chandorkar MS (1994) Highly conserved DNA sequence in *Isoetes pantii*. *Indian Fern J* 11: 53-5.
23. Goswami HK, Sharma US (1995) Large spores in Microsporangia of *Isoetes pantii* are megaspores. *Indian Fern J* 12: 195-6.
24. Goswami HK, Bhu I (1998) Bisexual sporangia in *Isoetes pantii*. *Bionature* 18: 31-6.
25. Goswami HK (2012) Directional selection of Bisexual plants in Natural populations of *Isoetes pantii* (Pteridophyta: Isoetaceae) with Chromosome Evolution. *Acta Bot Hungarica* 54: 103-15.
26. Goswami HK (2014) Heterosporangium: A New Category of Sporangium in Lycopsidea. *Acta Bot Hungarica* 56: 77-92.
27. Goswami HK (2011) Population cytogenetic studies confirm X-Y mechanism in *Isoetes pantii* (Isoetaceae: Pteridophyta). *Bionature* 31: 13-22.
28. Goswami HK, Arya BS (2012) *Isoetes* × *pantii* (Isoetaceae: Pteridophyta): A Review. *Nelumbo* 54: 187-92.
29. Verma SC (2018) Opportunities and challenges of exploring ferns and Lycophytes in the genomics age: A brief overview of some issues. In *Pteridology Today: Challenges and opportunities*. Edts: S C verma, S.P. Khullar , A. Benniamin P. Laxminarsimhan & Paramjit Singh Pp 8-32.
30. Goswami HK, Bhu I (2000) *Isoetes pantii* is a Natural hybrid: *I. sampathkumarini* x *I. coromandelina*. *Bionature* 20: 9-18.
31. Bajpai AK, Goswami HK (2002) Human Genome may share some genes with plant gene pool. *Bionature* 22: 63-5.
32. Bhu I, Goswami HK, Sharma US, Bajpai AK (2001) *Isoetes fuchsii*: A new *Isoetes* from India. *Bionature* 21: 11-7.
33. Rao LN (1944) New species of *Isoetes* from Bangalore, Mysore state. *Current Sci* 13: 286-7.
34. He Zican, Wang H, Li J, Ye Q, Taylor WC (2004) Chromosome behaviour during meiosis and development of spore mother cells in the Chinese quillwort *Isoetes sinensis* TC Palmer. *Amer F J* 94: 183-95.