

## Isolation of *Staphylococcus aureus* from Ice-Cream Samples

Samir H, Younis W, Sultan S and Abd El-Azeem MW\*

Faculty of Veterinary Medicine, Department of Microbiology, South Valley University, Qena, Egypt

\*Corresponding author: : Abd El-Azeem MW, Faculty of Veterinary Medicine, Department of Microbiology, South Valley University, Egypt, Tel: +201155842340, E-mail: wael\_2000uk@yahoo.co.uk

Citation: Samir H, Younis W, Sultan S and Abd El-Azeem MW (2018) Isolation of *Staphylococcus aureus* from Ice-Cream Samples. J Vet Ani Res 1: 204

### Abstract

Milk and dairy products including ice cream are good media for growth of *Staphylococci*, and dairy products are common sources of *Staphylococcal* intoxication. So the aim of this study is to detect the presence of *Staphylococcus aureus* in ice-cream samples which can be achieved by the following, 100 ice-cream samples were examined for presence of *S.aureus* on mannitol salt agar, the suspected colonies were identified by Gram staining and biochemically through catalase, oxidase and coagulase tests. Molecular identification of *S.aureus* by polymerase chain reaction (PCR) through detection of *16SrRNA* gene and *clfA* gene specific for *S.aureus*. Antimicrobial susceptibility testing were carried out through disc diffusion method to detect the susceptibility of all isolates to various antibiotics and molecular detection of *mecA* gene. This study showed that 22 out of 100 ice-cream samples were appearing positive *Staphylococci* through conventional methods of isolation. 15 samples were positive *S.aureus* according PCR technique. PCR detection of *mecA* gene showed that 10 out of 15 (66.6%) isolates have *mecA* gene. Disc diffusion method showed most of *Staphylococcus aureus* isolates were multi antibiotic resistant (MAR) to oxacilin, penicillin, rifampin and nalidixic acid but they were sensitive to chloramphenicol, vancomycin and tetracycline.

**Keywords:** Ice-Cream; *Staphylococci*; Antibiotic; *mecA* and *clfA*

### Introduction

Ice cream is one of the dairy desserts, it is a popular frozen food consumed particularly in summer, as well as, throughout all the year. It continues to present a dominant interest for a large segment of population. Many studies from different countries revealed that ice-cream also acts as a vehicle of food-borne diseases [1-3].

The *Staphylococci* are ubiquitous in nature, with humans and animals as the primary reservoirs. They are present in the nasal passages and throat, in the hair, and on the skin of probably 50% or more of healthy individuals. These organisms are associated with sore throats and colds, and are found in abundance in postnasal drip following colds. *Staphylococci* can be isolated from animals, with the bovine being the most important because of the involvement of *Staphylococci* in mastitis. Although animals and humans are the major source, *Staphylococci* also can be found in the air, dust, water, and human and animal wastes [4].

Milk and dairy products are excellent growth media for a large number of microorganisms, including *Staphylococci* [5]. Bacterial contamination of milk usually occurs during the milking process and this depends on the sanitary condition of the environment and utensils used for milking and the milker's hands, also it can gain access to milk by direct excretion from udders with clinical or subclinical *staphylococcal* mastitis [6,7].

*Staphylococcus* is one of the major bacterial pathogens which cause food poisoning [8]. *Staphylococcal* food poisoning (SFP) is a mild intoxication occurring after the ingestion of food containing *Staphylococcal* enterotoxins (SEs) [9]. There were five major classical SEs types, named; SEA, SEB, SEC, SED, and SEE. But now, new genes encoding enterotoxin such as SEG to SEU are identified. One or more of these genes are thought to be involved in *Staphylococcal* food poisoning [10].

Antimicrobial resistance is an important public health concern worldwide. The development of resistance both in human and animal bacterial pathogens has been associated with the extensive therapeutic use of antimicrobials or with their administration as growth promoters in animal production [11]. *Staphylococci* have been reported to frequently show multiple antimicrobial resistance patterns [12]. This may be due to the indiscriminate use of antibiotics has led to the development of multiple antibiotic resistances thereby rendering the antibiotic treatment ineffective [5]. The utilization of antibiotics in periods shorter than the recommended can also contribute to the antibiotic resistance.

Multiple antibiotic resistant *Staphylococcus aureus* (*S.aureus*) strains have been isolated from milk obtained from cattle, beef and human samples in many parts of the world [13, 14]. The prevalence of antibiotic resistance usually varies between isolates from the different sampled stations and even between isolates from different herds on the same farm [15]. There is many studies were evaluated the *S.aureus* in Ice cream, the difference between them and this manuscript was comparing the conventional and molecular methods in detection of *S.aureus*.

## Materials and Methods

### Sample collection

A total 100 samples of traditional ice cream were collected from different markets in Qena city with different flavor (17 with vanillia, 19 with chocolate, 13 with strawberry, 14 with mango, 14 with bananas and 23with mixed taste) within the shelf life period .The samples were transported to laboratory in sterile and cold containers (4 °C) and preserved at this temperature. The samples were processed immediately upon arrival using aseptic techniques [16].

### Isolation and identification of *Staphylococci* from ice cream

#### Isolation of *Staphylococci*

One gram of each sample was diluted with 9 ml of 1% buffered peptone water and homogenized in a stomacher for about 10 minutes [17].The diluted samples were plated onto mannitol salt agar (MSA). The plates were incubated aerobically at 37 °C for 18-24h. Characteristic *Staphylococci* colonies were further purified by sub-culturing onto MSA plates and the plates were incubated aerobically at 37 °C for 18 h–24 h. These isolates were retained for further bacterial identification [18].

#### Identification of the bacterial culture under microscope

Smears from the purified colonies were stained with Gram's stain and examined microscopically under oil immersion lens [19]. The typical colonies were showed gram-positive cocci occurring in bunched, grapelike irregular clusters were taken as presumptive *Staphylococcus* species.

#### Biochemical identification

Biochemical tests were performed to confirm *S.aureus* using Catalase test, Oxidase test and Coagulase test [20]. All *S.aureus* isolates were positive for catalase and coagulase and all of them were negative for oxidase test.

#### Molecular identification by PCR

In this study PCR technique was carried to confirm the presence of *S.aureus* in 22 ice-cream samples which selected by detection of *16S rRNA* gene specific for *Staphylococci* and *clfA* gene specific for *S.aureus* from culture samples Through the following steps [21].

#### DNA Isolation

It was carried out according to QIAamp DNA Mini Kit instructions (Catalogue no.51304).

Gene	Primer sequence (5'-3')	Length of amplified product	Reference	
<i>16S rRNA</i>	F-CCTATAAGACTGGGATAACTTCGGG	791 bp	(22)	
	R-CTTTGAGTTTCAACCTTGCGGTGCG			
<i>clfA</i>	F-GCAAAATCCAGCACACAGGAAACGA	638 bp		
	R-CTTGATCTCCAGCCATAATTGGTGG			
<i>mecA</i>	F-GTA GAA ATG ACT GAA CGT CCG ATA A	310bp		(24)
	R-CCA ATT CCA CAT TGT TTC GGT CTA A			

Table 1: Primers used for PCR

Gene	Primary denaturation	Secondary denaturation	Annealing	Extension	No. of cycles	Final extension
<i>16S rRNA</i>	94 °C 5 min.	94 °C 30 sec.	55 °C 40 sec.	72 °C 45 sec.	35	72 °C 10 min.
<i>clfA</i>	94 °C 5 min.	94 °C 30 sec.	55 °C 40 sec.	72 °C 45 sec.	35	72 °C 10 min.

Table 2: Cycling conditions of the primers during cPCR

**PCR Amplification:** For PCR amplification, using specific primers for each gene as shown in Table 1. A uniplex reaction mixture 25µl contained 1µl of Forward primers, 1µl of Reverse primers for each gene separately; 12.5µl Emerald Amp GT PCR master mix (2x premix), 6µl Template DNA and 4.5µl PCR grade water. The tubes were subjected to thermal cycling (Biometra) with programme

described as shown in Table 2. Amplified products were separated by agarose gel electrophoresis (1.5%) agarose containing 0.5 mg ethidium bromide per ml (sigma) Visualized and photographed under UV illumination. The sizes of the amplication products were estimated by comparison with a 100 bp DNA ladder (Fermentas. cat. no. SM0243).

### Antimicrobial susceptibility test

Antimicrobial susceptibility tests were performed by disc diffusion method according to the guidelines of Clinical and Laboratory Standard Institute [23]. Sensitivity pattern of the isolates were determined against Oxacillin (1mcg), Vancomycin (30mcg), Cefotaxime (30mcg), Chloramphenicol (30mcg), Nalidixic acid (30mcg), Rifampin (5mcg), Penicillin G (10mcg) and Tetracycline (30mcg). Antimicrobial testing results were recorded as sensitive, intermediate sensitive and resistant according to zone diameter interpretative standards provided by [23].

### Molecular detection of *mecA* gene by PCR

A uniplex reaction mixture 25µl contained 1µl of Forward primers 1µl of Reverse primers for each gene separately; 12.5µl Emerald Amp GT PCR master mix (2x premix), 6µl Template DNA and 4.5µl PCR grade water. The amplification was carried out with the following conditions: 94 °C for 5 min as Primary denaturation, 94 °C for 30 sec denaturation, annealing at 50 °C for 30 sec, extension at 72 °C for 30 sec and final extension at 72 °C for 7 min for 35 cycles.


## Results

### Isolation and identification of *Staphylococcus* isolates from ice cream samples

According to conventional method of identification through culture on mannitol salt agar, microscopic and biochemical identification, they were 22 samples positive for *S.aureus*. They have yellow colonies on mannitol salt agar; microscopically appear gram positive cocci, arranged in clusters, non-spore forming bacteria, positive catalase test and negative oxidase. Coagulase test showed that 4 isolates were strong coagulase and 8 isolates were suspected coagulase (weak coagulase).


### Molecular identification of *Staphylococci*

All 22 isolates were examined by PCR using *16S rRNA* gene specific for genus *Staphylococcus* and *clfA* gene specific for *S.aureus*. 15 out of 22 isolates were positive *Staphylococci* as shown in Figure 1, and all of them were *S.aureus* according *clfA* gene result as show in Figure 2.



Lane L: DNA ladder, Lane pos: control positive *Saphylococci*, Lane neg: control negative, Lane 1,2,3,4,5,6,7,8,9,10,11,13,14,15 positive *Staphylococcus* isolates, Lane 12,15,17,18 negative isolates.

**Figure 1:** PCR results for the *16S rRNA* gene (791bp)



Lane L: DNA ladder, Lane pos: control positive *S.aureus*, Lane neg: control negative, Lane 1,2,3,4,5,6,7,8,9,10,11,13,14,15,22 positive *S.aureus* isolates

**Figure 2:** PCR results for *clfA* gene (638bp)

## Results of antimicrobial susceptibility testing


All the 15 isolates were examined for antibiotic sensitivity to different antibiotics through disc diffusion method. Antimicrobial susceptibility testing through disc diffusion method shows the following (Table 3):

Antibiotic disc	Sensitive		Intermediate		Resistant	
	No	%	No	%	No	%
Penicillin (p)	1	6.6%	-	-	14	93.3%
Oxacillin (OX)	1	6.6%	2	13%	12	80%
Chloramphenicol (C)	14	93.3%	-	-	1	6.6%
Tetracycline (TE)	11	73.3%	-	-	4	26.6%
Vancomycin (VA)	12	80%	2	13%	1	6.6%
Cefotaxime (CTX)	8	53.3%	6	40%	1	6.6%
Rifampin (RA)	5	33.3%	2	13%	8	53.3%
Nalidixic acid (NA)	6	40%	-	-	9	60%

**Table 3:** Results of sensitivity of different *S.aureus* isolates to different antibiotic discs

## Occurrence of *mecA* gene among *S.aureus* isolates

Among the 15 *S.aureus* strain, there were 10 strains have *mecA* gene, as show in Figure 3.



Lane L: DNA ladder, Lane pos: control positive *mecA*, Lane neg: control negative, Lane 2,5,6,7,8,10,11,14,15,22 (+*mecA*), Lane 1,3,4,13 (-*mecA*)

**Figure 3:** PCR results of *mecA* gene (310bp) among *S.aureus*

## Discussion

Ice cream is one of the most abundant and popular dairy products that are consumed in warm seasons by vulnerable groups, especially children, therefore; its microbial contamination is very important. This study has shown contamination of ice-cream with *S.aureus*.

The present study aimed to detect *S.aureus* and detection of antibiotic resistance profile and methicillin resistance gene of *S.aureus* isolated from ice cream.

A total 22 (22%) *S.aureus* isolates were detected from 100 samples of ice cream, all the isolates were gram positive cocci arranged in clusters, catalase positive and oxidase negative which agree with who isolated *S.aureus* from 11/50 (22%) ice-cream samples. And nearly similar finding to that assumed by who isolated *S.aureus* from 26%, 22.9% and 20%, respectively [25-28].

On the other hand, higher incidence was reported by who isolated *S.aureus* from 84.72%, 50%, 76%, 56.67% and 55% respectively, but lower incidence was reported by isolated *S.aureus* from 2.7 %, 4.3%, 0.5%, 12.2% and 4.4%, respectively. Moreover, could not detect *S.aureus* in any one of the examined ice cream samples. The difference in the incidence may be due to bad hygienic measurement during manufacture of ice cream [29-42].

In this study the 22 *S.aureus* isolates were conducted for PCR and we found 15 out 22 (68.2%) showing positive using *16SrRNA* gene and all of them were *S.aureus* using *clfA* gene specific for *S.aureus* which in agreement with because the PCR is highly sensitive while conventional method is less sensitive as there is some microorganism give positive reaction by culture and biochemical tests but give negative by PCR [21].



Antimicrobial resistance is an important public health concern worldwide. It has been believed that all bacterial infections treated with effective antimicrobial agents. However, the emergence of resistance to multiple antibiotics among *S.aureus* has created breaking news for health practitioners and researchers [43]. It has been reported shortly after introduction of penicillin 1940s, resistance developed in *S.aureus* followed by resistance to methicillin and more recently to glycopeptides as vancomycin [44].

In this study the isolates were evaluated for antimicrobial resistance through disk diffusion method, resistance to penicillin G was high (93.3%), high resistance to  $\beta$ -l actam antibiotic was not surprising as it is commonly used for treatment of infections in humans and animals [45].

The present study *S.aureus* was resistant to oxacillin (80%), this result nearly similar to and who found resistance of *S.aureus* to oxacillin 76.2% and 86.7% respectively. While, this result disagreed with that of and who detected resistance rate against oxacillin of 6.2% and 28% respectively [46-49].

This study show that one *S.aureus* isolate sensitive to oxacillin and contain *mecA* gene which nearly similar to who obtained results demonstrated low correlation ( $p>0.05$ ) between phenotypic resistance to oxacillin and the presence of *mecA* gene in *Staphylococci* and found the similar results in phenotypically oxacillin resistant isolates of *S. aureus*. Those strains did not carry neither *mecA* nor *mecC* genes [50,51].

Furthermore, there were four isolates show oxacillin resistance phenotypically and could not have *mecA* gene which agree with who found only 5 isolates have *mecA* gene out of 22 isolates showing oxacillin resistance phenotypically by disc diffusion method. Oxacillin has been proposed as a proxy antibiotic for testing susceptibility not only to methicillin and to all  $\beta$ -lactams, which could explain why all oxacillin-resistant isolates were not carrying the *mecA* gene [52,53].

The present study shows that *S.aureus* isolates were resistant to vancomycin 6.6%. This nearly agree with who found *Staphylococcus* isolates were resistant to vancomycin with 3.33% and disagree with who reported that *S.aureus* were 100% resistant to vancomycin [54,55].

The present study show that *S. aureus* isolates were sensitive to chloramphenicol 93.3% which nearly agree with who found 85.96% of *S.aureus* isolates sensitive to chloramphenicol and who reported that all *S.aureus* isolates were susceptible to chloramphenicol. These results were not in agreement with and who found resistance of *S.aureus* isolates to chloramphenicol 33.3% and 25% respectively. Also, it disagree with who found that *S.aureus* were resistant to chloramphenicol in 80% [46, 55-57].

The resistance to tetracyclines occurs through the presence of *tet* genes in the bacterial DNA. The characterized *tet* genes encode three mechanisms of resistance: efflux pump, ribosomal protection or enzymatic inactivation 28 [59].

In this study, 73.3% of the isolates were susceptible to tetracycline which nearly agree with who found about 75.44% of the isolates were sensitive to tetracycline. And disagree with and who reported that *S.aureus* were resistant to tetracycline in 74.1% and 85% respectively. Also it disagrees with who observed that all isolates of *S.aureus* (100%) were resistant to tetracycline [4, 46,56,58].

About 53.3% of the isolates were resistant to rifampin which were disagree with who reported that 68.4% of *S.aureus* isolates were sensitive to rifampin [56].

The sensitivity of *S.aureus* isolates to cefotaxime is (53.3%) which were not in agreement with who reported that, all *S.aureus* isolates were susceptible to cefotaxime [57].

In this study, the isolates were resistant to nalidixic acid in 60% which were not in agreement with who found that all *S.aureus* isolates 100% were resistant to nalidixic acid. The difference in resistance of *Staphylococcus aureus* isolates to different antibiotics may be due to genetic variation and phenotypic variation.

It was observed that 66.6% of *S.aureus* isolates were positive for presence of *mecA* gene and this in agreement with who found 66.7% of *S.aureus* isolates were positive for presence of *mecA* gene and who isolate MRSA in 50% and disagree with and who detect MRSA in 18.18% and 29.6% respectively [46, 61-63].

## Conclusion

Ice cream is one of most popular and favorite food products all over the world. It is an ideal media for microbial growth due to high nutritive value and long storage duration. Once the ice becomes cream contaminated, freezing temperature could not make the product safer later.

This study revealed that, some ice cream sold in Qena city, Egypt was contaminated with *Staph. aureus* which may cause food poisoning. It indicates the lack of hygienic conditions during preparation and preservation of ice cream. Antimicrobial resistance is an important public health concern worldwide. In the present study, *Staph.aureus* isolates exhibit resistance towards the different antibiotics tested.

Thorough food inspection and frequent bacteriological surveillance by food control agencies is highly recommended to control the incidence of *Staph.aureus* in dairy products to safeguard the consumers from risks of food poisoning.

## References

1. Hussein MF, Sadek OA, El taher SG (2015) Occurrence of bacillus cereus and *Staphylococcus aureus* organisms in some dairy desserts. Assiut Vet Med J 61: 145.
2. Seidel G, Janeck K, Hoffman G (1962) An occurrence of *Staphylococcus aureus* food poisoning caused by ice-cream. Mh Vet Med 17: 782.
3. Dardzinski K (1965) Clinical and epidemiologic analysis of mass *Staphylococcal* food poisoning caused by ice-cream. Przegł Epidem 19: 245.
4. Bergdoll M, Lee WA (2006) *Staphylococcal* intoxications. In: Foodborne Infections and Intoxications; Academic Press, Elsevier: USA: 523–25.
5. Farzana K, Hussain Shah SN, Jabeen F (2004) Antibiotic resistance pattern agent various isolates of *Staphylococcus aureus* from raw milk samples. J Res 15: 145–51.
6. Smith K, Peter K, Daniela H, Melchior S (2007) Food borne pathogenic microorganisms and natural toxins. Food drug Administration Center Food Safety Applied Nutrition 10: 119–50.
7. Peles F, Wagner M, Varga L, Hein I, Rieck P, et al. (2007) Characterization of *Staphylococcus aureus* strains isolated from bovine milk in Hungary. Int J Food Microbiol 118: 186–93.
8. Chiang YC, Liao WW, Fan CM, Pai WY, Chiou CS, et al. (2008) PCR detection of *Staphylococcus enterotoxins* N, O, P,Q,R,U and survey of SE types in *S. aureus* isolates from food poisoning cases in Taiwan. Int J Food Microbiol 121: 66–73.
9. Chiang YC, Liao WW, Fan CM, Pai WY, Chiou CS, et al. (2008) PCR detection of *Staphylococcus enterotoxins* N, O, P,Q,R,U and survey of SE types in *S. aureus* isolates from food poisoning cases in Taiwan. Int J Food Microbiol 121: 66–73.
10. Afifi AN, Sadek A, Aggour GM, El-Temawy AA, Nemr MM (2011) Molecular Characterization of *Staphylococcus aureus* Enterotoxins in Milk and Some dairy Products. Egypt J Med Microbiol 20: 107–16.
11. Normanno G, La Salandra G, Dambrosio A, Quaglia NC, Corrente M, et al. (2007) Occurrence, characterization and antimicrobial resistance of enterotoxigenic *Staphylococcus aureus* isolated from meat and dairy products. Int J Food Microbiol 115: 290–6.
12. Enright MC (2003) The evolution of resistant pathogen - the case of MRSA. Current Opinion in Pharmacol 3: 474–9.
13. Shitandi A, Sternesjö Å (2004) Prevalence of multidrug resistant *Staphylococcus aureus* in milk from large and small-scale producers in Kenya. J Dairy Sci 87: 4145–9.
14. Petinaki E, Miriagou V, Hatzi F, Kontos F, Maniati M, et al. (2001) Bacterial Resistance Study Group. Survey of methicillin-resistant coagulase-negative *Staphylococcus aureus* in the hospitals of central Greece. Int J Antimicrob Agents 18: 563–6.
15. Waage S, Bjorland J, Caugant DA, Oppegard H, Tollersrud T, et al. (2002) Spread of *Staphylococcus aureus* resistant to penicillin and tetracycline within and between dairy herds. Epidemiol Infect 129: 193–202.
16. Mirzaei H, Farhoudi H, Tavasoli H, Farajli M, Monadi A (2012) Presence and antimicrobial susceptibility of methicillin resistant *Staphylococcus aureus* in raw and pasteurized milk and ice cream in Tabriz by culture and PCR techniques. Afr J Microbiol Res 6: 6224–9.
17. Gundogan N, Atal O (2013) Biofilm, protease and lipase properties and antibiotic resistance profiles of *Staphylococci* isolated from various foods. Afr J Microbiol Res 7: 3582–8.
18. Daka D, silassie SG, and Yihdego D (2012) Antibiotic-resistance *Staphylococcus aureus* isolated from cow's milk in the Hawassa area, South Ethiopia. Ann Clin Microbiol and Antimicrobials 11: 26.
19. Quinin PJ, Carter ME, Markey BK, Donnoly WJ, Leonard FE (2002) Veterinary microbiology and microbial disease. 166-1117 Osney Mead Oxford first LTD Registered at the United Kingdom.
20. Sneath PHA, Mair NS, Sharpe ME, Holt JG, Murray RGE, et al. (1986) Section 12 gram positive cocci, Bergey's manual of systemic bacteriology, Williams & Wilkins USA.
21. El-Nagar S, Abd El-Azeem MW, Nasef SA, Sultan S (2017) Prevalence of Toxigenic and Methicillin Resistant Staphylococci in Poultry Chain Production. J Adv Vet Res 7: 33–8.
22. Mason WJ, Blevins JS, Beenken K, Wibowo N, Ojha N, et al. (2001) Multiplex PCR Protocol for the Diagnosis of *Staphylococcal* Infection. J Clin Microbiol 39: 3332–8.
23. CLSI (Clinical and Laboratory Standards Institute) (2011) Performance Standards for Antimicrobial Susceptibility Testing; Twenty- First Informational Supplement. M100-S21 30: 15.
24. McClure JA, Conly JM, Lau V, Elsayed S, Louie T, et al. (2006) Novel multiplex PCR assay for detection of the *Staphylococcal* virulence marker Pantone-Valentine leukocidin genes and simultaneous discrimination of methicillin-susceptible from -resistant *Staphylococci*. J Clin Microbiol 44: 1141–114.
25. Abd El, Tawab AA, Ammar MA, El-Hofy IF, Aideia AH, et al. (2016) Bacteriological and molecular studies on toxigenic *Staphylococcus aureus* in milk and some milk products. BVMJ 31: 202–9.
26. Abdel-Fatah EN (2010) Role of small and large scale producers in transmission of food poisoning organisms through consumption of ice-cream. Ph.D. Thesis, Zag Univ, Egypt.
27. Kamal RM (2009) Sanitary status of some locally and imported dairy products. Ph.D. Thesis, Zag Univ, Egypt.
28. Masud T (1989) Microbiological quality and public health significance of ice-cream J Pak Med Assoc 39: 102–4.
29. Abdel-Haleem AA (1995) Microbiological evaluation and sanitary improvement of ice-cream. Ph. D. Thesis, Assiut Univ Egypt.
30. Abo-Risha NE. (1998) Occurrence of some food poisoning microorganisms in some dairy products. M.V.Sc. Thesis, Kafr El-Sheikh Univ, Egypt.
31. Allam HA (1999) Microbiological studies on milk and some milk products. Ph.D. Thesis, Benha Univ, Egypt.
32. Hammad AM (2004) Microbiological studies on raw milk and some dairy products M.V.Sc., Thesis, Minufiya Univ, Egypt.
33. Hassan GM (2003) Quality assessment of some dairy products at consumer level .M.V.Sc., Cairo Univ, Egypt.
34. Amurajimi C, Geethu S, Dhanashree B (2008) Bacteriological analysis of ice-cream from Mangalore. South India. Indian J Med Res 127: 91–2.

35. Kock C, Akan M, Yardimci H (1998) Bacteriological quality of ice-cream marketed in Ankara. *Veterinary-Fakultesi-Dergisi-Ankara Univ* 45: 113-34.
36. Kruiy SL, Soares JL, Ping S, Sainte-Marie FF (2001) Microbiological quality of “ ice cream.Sorbet “ sold on the streets of Phnom Penh; April 1996- April 1997. *Bull Soc Pathol Exot* 94: 411-21.
37. Little CL, Louvois J (1999) Microbiological Quality of soft ice-cream from fixed premises and mobile vendors. *Int J Environ Health Res* 9: 223-32.
38. Manzanera-Pelegrin C, Marin-Mesegure D, Paredes-Garcia P, Pelegrin CM, Mesegure DM, et al. (1995) Quality control of ice-cream. *Health area III of Murcia region Alimentaria* 33: 119-21.
39. Caglayanlar GE, Buket Kunduhoglu B, Coksoyler N (2009) Comparison of microbiological quality of packed and unpacked ice-cream sold in Bursa. *Turk J Arts Sci* 12: 93-102.
40. Korel F, Omeroglu S, Tan G, Odabaso AZ, (2002) The evaluation of chemical and microbiological quality of ice-cream sold in retail markets in Manisa. *Turk. Annual Meeting and food Expo- Anaheim, California*.
41. Maiereni M, Civilini M, Domenis C, Manzano M, Di-Prima R, et al. (1993) Microbiological quality of Artisanl ice cream. *Zentralbl Hyg Umwelt Med Bio Abst* 194: 553-70.
42. Sagdic O, Tuluoglu DD, Ozcelik S, Simsek B (2002) The chemical and microbiological quality of ice-cream consumed in Isparta market Ziraat. *Fakultesi-Dergisi, Ataturk Uni* 33: 441-6.
43. Hossein M, Hadis M, Tahere S (2010) Determining of antibiotic resistance profile in *Staphylococcus aureus* isolates. *Asian Pac J Trop Med* 3: 734-77.
44. Monroe S, Polk R (2000) Antimicrobial use and bacterial resistance. *Curr Opin Microbiol* 3: 496-501.
45. Gundogan N, Citak S, Yucel N, Devren AA (2005) note on the incidence and the antibiotic resistance of *Staphylococcus aureus* isolated from meat and chicken samples. *Meat Sci* 69: 807-10.
46. Ibraheem TES (2017) Molecular characterization of toxigenic *Staphylococcus* isolated from chicken production cycle. Ph.D. thesis, S V U, Egypt.
47. Eid S, Nasef SA, Erfan AM, (2015) multidrug resistant bacterial pathogens collected from backyard chickens. *Assuit Vet Med J* 61: 144.
48. Nam HM, Lee AL, Jung SC, Kim MN, Jang GC, et al. (2011) Antimicrobial susceptibility of *Staphylococcus aureus* and characterization of methicillin- resistant *Staphylococcus aureus* isolated from bovine mastitis in Korea. *Food borne pathog Dis* 8: 231-8.
49. Giedam YA, Zakaria Z, Abdul Aziz S, Bejo SK, Abu J, et al. (2012) High prevalence of multidrug resistant bacteria in selected poultry farms in Selanger, Malaysia. *Asian J Anim Vet Adv* 7: 891-7.
50. Ružauskas M, Vaskeviciūtė, Šiugždinienė R, Klimienė I, Virgaillis M et al. (2014) Antimicrobial susceptibility of oxacillin-resistant *Staphylococcus* Spp. Isolated from poultry products. *Microbiology and Virology Institute, Lithuanian University of Health Sciences*.
51. Ba X, Harrison EM, Edwards GF, Holden MTG, Larsen AR, et al. (2014) Novel mutations in penicillin –binding protein genes in clinical *Staphylococcus aureus* isolates that are methicillin resistant on susceptibility testing, but lack the mec genes. *J Antimicrob Chemother* 69: 594-7.
52. Akanbi EO, Njom AH, Frij, Otiqbu CA, Clarke MA (2017) Antimicrobial Susceptibility of *Staphylococcus aureus* Isolated from Recreational Waters and Beach Sand in Eastern Cape Province of South Africa. *Int J Environ Res Public Health* 14: 1001.
53. Kuehnert MJ, Hill HA, Kupronis BA, Tokars JL, Solomon SL, et al. (2005) Methicillin-resistant- *Staphylococcus aureus* hospitalizations, United States *Emerg Infect Dis* 11: 468-872.
54. Piyali D, Pranab BM (2016) Prevalence of *Staphylococcus* in raw meat samples in Southern Assam, India, *J Agric Vet Sci* 9: 23-9.
55. Shareef AM, Mansour RS, Ibrahim, KK (2009) *Staphylococcus aureus* in commercial breeder layer flocks. *Iraqi J Vet Sci* 23: 63-8.
56. Ghadimi S, Heshmati A, Azizi, Shafa M, Nooshkam M (2017) Microbial Quality and Antimicrobial Resistance of *Staphylococcus aureus* and *Escherichia coli* Isolated from Traditional Ice Cream in Hamadan City, West of Iran. *Avicenna J Clin Microb Infec* 4: e39781.
57. Gundogan N, Avaci E (2014) Occurrence and antibiotic resistance of *Escherichia coli*, *Staphylococcus aureus* and *Bacillus cereus* in raw milk and dairy products in Turkey. *Int J Dairy Technol* 67: 562-9.
58. Yurdakul NE, Erginkaya Z, Unal E (2013) Antibiotic resistance of enterococci, coagulase negative and *Staphylococcus aureus* isolated from chicken meat. *Czech J Food Sci* 31: 14-9.
59. Sumru C, Tugba D (2011) *Staphylococcus aureus* and coagulase negative *Staphylococcus* from raw chicken samples in Turkey: prevalence and antimicrobial resistance, *Food, Agriculture and Environment* 9: 156-8.
60. Chigbu C, Ezeronye OU (2003) Antibiotic resistant *Staphylococcus aureus* in Abia State of Nigeria. *Afr J Biotechnol* 2: 374-8.
61. Febler AT, Wendlandt S, Ruppelt A, Kaddlee K, Hassel M, et al. (2012) Methicillin- resistant *Staphylococcus aureus* (MRSA) from poultry and food of poultry origin: molecular characterization and antimicrobial resistance- Brazil 5-9.
62. Akbar A, Anal K (2013) Prevalence and antibiogram study of salmonella and *Staphylococcus aureus* in poultry meat. *Asian Pac J Trop Biomed* 3: 163-8.
63. El-Shareek, YM, Ali, MRM (2012) Microbiological study of spiced chicken burgers in Tripoli City, Libya. *EMHJ* 18: 653-62.