

Physiochemical and Functional properties of (*Balanites aegyptiaca. del*) hydrolysates by pepsin and pancreatin proteases.

Ogori AF^{1,*}, Eke MO², Girgih TA² and Abu JO²

¹Faculty of Agriculture, Department of Home Sciences Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria

²College of Food and Human Ecology, Department of Food Science and Technology, Joseph Sarwuan Tarka University, Makurdi, Benue State, Nigeria

Corresponding Author: Akama Friday Ogori, Faculty of Agriculture, Department of Home Sciences Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria, Tel: 2348070626328, E-mail: ogorifaraday@gmail.com.

Citation: Akama Friday Ogori, Eke MO, Girgih TA and Abu JO (2023) Physiochemical and Functional properties of (*Balanites aegyptiaca. del*) hydrolysates by pepsin and pancreatin proteases. J Food Technol Food Chem 5: 101

Abstract

Enzymatic hydrolysates from underutilized desert date could produce functional ingredient for desert region and population of the world. The current study investigated the physiochemical and functional properties of hydrolysates from *Balanites aegyptiaca* using pancreatin, pepsin and combined pancreatin and pepsin enzymes. The combined hydrolysate had (45.1%) yield. On the other hand, pancreatin hydrolysate gave maximum (45.9%) yield and pepsin hydrolysate was (37.4%). The combined hydrolysate has maximum protein (80.58%), ash content (0.85%) mineral profile; Ca(0.12mg/100g), K(1.29mg/100g), Na(2.01mg/100g), Mg (1.09mg/100g), Fe(0.23mg/100g), Cu (0.09mg/100g) , Zn(0.26mg/100g) and excellent anti-nutrient alkaloid , oxalate , phytate and saponin and total phenolic content of (0.084mgCat/g). Functional properties such as bulk density (1.02g/m) swelling index (3.45g/m) least gelation capacity (20.39%) of pancreatin hydrolysate and WAC (1.72g/g) and OAC (1.73g/g) from pepsin hydrolysate produced the best result than the combine hydrolysate. The maximum solubility (12%) at pH 9 was observed for all samples. The hydrolysate by combined enzymes had a high IVPD compared with pancreatin and pepsin hydrolysates. However, both showed a good general foaming and emulsifying properties. Foaming capacity of the pepsin hydrolysate samples increased with concentration at pH3 and pH7 while the foaming stability at pH7 and pH9 of combined hydrolysate was at 20 mg/ml and 40 mg/ml high comparable with pepsin and pancreatin, respectively. The emulsifying activity of combined hydrolysate was high at 10mg/ml and 50mg/ml at alkaline region. On the other hand, pepsin hydrolysate reveals excellent emulsifying activity (EAI) of 200 mg/m index (EAI) at pH 5. The hydrolysate obtained from pepsin decreased in emulsifying stability as concentration increased and pH shift to alkaline region. The hydrolysate by pepsin and pancreatin had better WHC, OHC and foaming properties due to their solubility however lower than the combined hydrolysate. The result indicates potential utilization of hydrolysate from *Balanites aegyptiaca. del* for less protein rich cereal food formulations.

Key word: Physical; Chemical; Functional; Pepsin; Pancreatin; Hydrolysate

Introduction

The need to exploit more proteins and peptides from diverse natural resources are on the increase and with respect to climate change, it may double soon. A lot of forest and desert reserved plant source of food abound and needs to be exploited for human maximum beneficial use. Protease hydrolysis is an excellent way for solubilizing and exposing peptides conformation for cellular utilization of their essential amino acids or proteins in living cells. Hydrolysis of protein is adoptable in small and industrial scale, widely used in the food industry to make stable or semi stable products which could serve as finished or raw material in milk as replacer, beverage stabilizer and flavor enhancer. Plant food hydrolysates when derived by protease hydrolysis, tends to have better nutritional profile in terms of amino acids, peptides classes which could be used for human or animal feeds [30]. The greater benefit of using plant source food as functional modifier for sourced free amino acid, reduce phytotoxins and give bioactives have been opined, [2,3]. Derivable proteins or peptides from hydrolysis are crucial in their use as food ingredient [4] and as potential therapeutic food or drug incipient. Enzymatic hydrolysis of plant protein using proteases such as pepsin, pancreatin Alcalase, Flavoenzyme and Chymotrypsin have been opined [34]. Food functions are related their physiochemical properties like amino acid profile, molecular weight, charge distribution [3]. It has been elaborated that hydrolysate stereochemistry relates with food biomolecules determined by intrinsic physical parameters are food product modifiers [3,6,7]. Aduwa seed have been reported utilizable for human and feed use, the use of Aduwa leaves, fruits and seed have also been researched and reported with potential utilization at homes, industries and pharmaceutical applications as anti-diabetes remedy, anti-cancer, anti- helminthics as well as an antioxidant [2,8]. The goal of this study is to examine the properties of desert date enzymatically hydrolyzed with pepsin and pancreatin from toasted Aduwa seed to direct their application and use towards food and pharmaceutical products for human use.

Material and Methods

Material

Aduwa seeds were cracked and packaged from bade in yobe state, Northeast of Nigeria. They were identified at forestry department and given lot number (Fror 23). The seed were toasted and milled before defatted. The defatted meal was extracted by isoelectric precipitation into concentrates and well packaged and ready for evaluation. Samples were taken and made into hydrolysate using pepsin and pancreatin at 4% w/v enzyme addition.

Making Aduwa Concentrate (APC)

The concentrates (AC) were prepared as described and modified [9]. About 200 g weighed defatted Aduwa meals were dissolved to final 1:10w/v ration. Slurry mixtures were stirred until a homogeneous mix before pH adjustment to pH 4. The precipitation process was allowed for 2h at constant pH 4 and stirring. The mix was centrifugation at $3,500 \times g$ for 30 min using centrifuge and the precipitate (concentrates) were washed thoroughly and then the pH was adjusted to pH7.0. The concentrates were collected and oven dried at 45 °C for 8 h and kept in air-tight container.

making Aduwa hydrolysate with pancreatin enzyme

Aduwa hydro lysate (APH) using pancreatin was prepared using the method of [10]. A 1:20 w/v Aduwa concentrate slurry was incubated at 40°C, adjusted to pH 7.5 following the addition of pancreatin at (4% w/w), based on protein content of Aduwa concentrate. The digestion was allowed for 4 h at constant pH. The digestion was stopped after adjusting the pH to 4.0. The mixtures were placed in a boiling water for 30 min for complete enzyme denaturation, cool to room temperature and later centrifuge for the supernatant and freeze dried.

Preparation of Aduwa protein hydrolysate using pepsin enzyme

The Aduwa meal protein hydrolysate (AMPH) was prepared using pepsin enzymes in an optimum reaction condition (Pepsin with pH 2 at 37°C), using the described method of [10]. A 1:20 w/v Aduwa protein concentrate slurry was adjusted to pH 2.0 and incubated at 37 °C followed by addition of pepsin (4% w/w based on protein content of Aduwa protein isolate. The digestion was carried out for 4 h and the pH is maintained by adding 1 M NaOH or HCl where necessary. The digestion was terminated by adjusting the pH to 4.0 and then place the mixtures in boiling water for 30 min to inactivate the enzymes. The mixture was allowed to cool to room temperature and later centrifuged and supernatant collected and freeze dried.

Enzyme hydrolysis of APC for making combined Hydrolysate (APHpa+pe)

Enzymatic hydrolysis of APC was carried out using the method of Aluko and McIntosh (2004) with slight modification by [4]. The APC sample was dispersed in water (2%, w/v), and adjusted to pH 9.0 using 1 M NaOH solution for pancreatin while pH 2.0 was used for pepsin digestion. The dispersion was heated to 40°C under continuous stirring on a hotplate. The enzymes (4% w/w) were added based on the protein content of the APC and incubated at constant temperature of 40°C for 4 h. The reaction mixture was maintained at pH 9.0 using 1 M NaOH solution and after 4 h, the pH was adjusted to pH 2.0 with 1 M HCl for pepsin hydrolysis. At the end of the incubation period, the hydrolysates were transferred into a boiling water bath for 5 min to inactivate the enzymes. The hydrolysate was cooled to room temperature (22±2 °C) using ice blocks and adjusted to pH 7.0 and finally freeze-dried.

Moisture content determination

Five grams (5g) of the hydro lysate sample was weighed and heated in a pre-weight crucible at 103°C for 3 hours. At the end, the dish was desiccated and allowed to cool for 30 minutes. The cooled dish was weighed and oven kiln again for another 2 hours until decreases in weight- mass was observed. Weight loss was reported thus.

$$\%Moisturecontent = \frac{W1 - W2}{W1 - W0} \times 100$$

Protein determination by Lowry Method

The [2] method was adopted. Here, one mg/ml of the soluble filtrate was pipetted with the addition of 3 ml of Lowry's reagent C was used and dissolved and make to the mark with distilled water in a 100 mL standard flask; Reagent X: (4 % CuSO₄.5H₂O) was dissolved and made-up to the mark with distilled water in 100 mL standard flask. The mixture was incubated at room temperature for 1 h. Also, 0.3 mL of diluted Folin Ciocateu phenol was added to the mixture and mixed vigorously using vortex mixer. The tubes were allowed to stand at room temperature for 45 min and the absorbance of the mixture was then measured at 600 nm using spectrophotometer. Bovine Serum Albumin (standard) was prepared in similar manner as the samples but at different concentration (1-100µg/mL). The standard curve obtained was used to find the protein concentration of the sample.

Ash content determination

Two grams (2g) of the hydrolysate sample was weighed into an empty porcelain crucible that was ignited and weighed. The hydrolysate sample was ignited over a hot plate in a fume cupboard to char. The crucible was thereafter placed in the muffle furnace maintained at a temperature of 600°C for 6 hr. After ash, samples were then transferred directly to a desiccator and weighed immediately [13].

$$Ash(\%) = \frac{(weight\ of\ the\ crucible\ and\ Ash) - (weight\ of\ empty\ crucible) \times 100}{weight\ of\ hydrolysate\ sample}$$

Crude fat determination

Crude fat determination was carried out using the method of [11]. Empty thimble was weight and recorded as W_1 . Five (5) grams of oven dried hydrolysate sample was added and weighed as (W_2). Round bottom flask was used in the Soxhlet extraction with petroleum ether as extracting solvent. Soxhlet extractor was fixed with a reflux condenser to adjust the heat sources so that the solvent boils gently. The samples were put inside the thimble and inserted into the Soxhlet apparatus and extraction under reflux was carried out with petroleum ether for 6 hours. After the barrel of the extractor became empty, the condenser and the thimble were removed. The thimble was taken to the oven at 100°C for 1 hour and later cooled in the desiccator. The sample was then weighed as (W_3).

$$\%Fat = \frac{\text{weight loss of extracted fat}}{\text{original weight of hydrolysate sample}} \times 100$$

Crude fiber determination

Crude fiber content of the samples was determined as described in [11]. Two (2) grams of sample was weighed into a 250ml volume beaker, allowed to boiled for 30 minutes when 100ml of 0.12M H_2SO_4 were added. The mix was filtered and the filtrate washed with boiling water repeatedly. The solution was boiled for another 30 minutes with 100ml of 0.012M NaOH solution; filtered and washed with water and alcohol. The final residue was put in a crucible and dried in the oven at 103°C for 1 h, cooled in a desiccator and then reweighed (W_1), before ash at 600°C for 1 h. The ashed sample was desiccated and weighed as (W_2). The percentage crude fiber was calculated thus:

$$\%Crude\ fiber = \frac{w_1 - w_2}{\text{weight of sample}} \times 100$$

Carbohydrate content determination

Carbohydrate content determination was determined by difference [14]

$$\%Carbohydrates = 100\% - \%(\text{Protein} + \text{Crude fat} + \text{Ash} + \text{Moisture} + \text{Crude fiber})$$

Tannins

The modified vanillin – hydrochloric acid (MV – HCl) method of [15] was used. Entails the preparation of standard solution and stock solution. The Calibration curve concentrations (0.0, 0.4, 0.8 and 1.0 mg/ml) of the catechin standard solution were pipetted into clean dried test tubes. A freshly prepared vanillin – HCl reagent was prepared by mixing equal volume of 4% (w/v) vanillin/MeOH and 8% (v/v) HCl/ MeOH added. The second set, 5.0 ml of 4% (v/v) HCl and methanol was Mix. This serve as blank. The solutions were left for 20 min before the absorbance was taken at 500 nm. The absorbance of the blank will be subtracted from that of the standards. Sample readings followed these extraction steps. The extraction time is 1 hour with continuous shaking, after shaking the solution was filtered and made up to 10 ml mark with extracting solvent. The filtrate (1.0 ml) was then reacted with 5.0 ml vanillin – HCl reagent and another with 5.0 ml of 4% (v/v) HCl – MeOH solution to serve as blank. The mix was left to stand for 20 min before the absorbance was taken at 500 nm. Tannin was calculated using the formular.

$$Tannin = \frac{X\text{mg/ml} \times 10\text{ml}}{0.2\text{g}} = 50 \times \text{mg/g}$$

Oxalate

Oxalate was determined by the method of [16]. Four grams of the sample were weighed in triplicate into 250 ml conical flasks and was extracted with distilled water and 6M HCl. The extract were allowed to boil for 2 h, filtered and made up to 250 ml with water. To 50 ml aliquot, 10 ml 6M HCl was added and filtered, and the precipitates were washed with 10 ml of hot water. The filtrate and the wash water were combined and titrated against concentrated NH_4OH until the salmon pink color of the methyl red indicator changed to faint yellow. The solution was heated to 90 °C and 10 ml 5 % (w/v) and CaCl_2 solution added to precipitate the oxalate overnight. The precipitates were washed free of calcium with distilled water into 100 ml conical flask with 10 ml hot 25% (v/v) H_2SO_4 and with 15 ml distilled water then heated to 90 °C and titrated against a standard 0.05M KMnO_4 until a faint purple solution persisted for 30 s. The oxalate was calculated as the sodium oxalate equivalent

Determination of saponin

The spectrophotometric method [17] was used for saponin analysis. One gram of finely ground hydrolystates samples was weighed into 250 ml beaker and 100 ml of isobutyl alcohol added. The mixture was shaken on a Brunswick incubator shake for 5 h. Thereafter the mixture was filtered through into a 250 ml beaker and 20 ml of 40% saturated solution of magnesium carbonate added and the mixture made up to 250 ml. The mixture that was obtained with saturated MgCO_3 was filtered to obtain a clear colorless solution. 1 ml of the colorless solution was pipetted into a 50 ml volumetric flask and 2 mL of 5% FeCl_3 solution added and made up to mark with distilled water. The mixture was allowed to stand for 30 min for blood red color to develop. Saponin stock was prepared at 0 – 10 ppm from saponin stock solution. The standard solution was treated similarly with 5% of FeCl_3 solution. The absorbance values of the sample as well as the standard solution was read after color development in spectrophotometer (722-2000 spectronic 20D, England) at a wavelength of 380 nm.

Phytate

The phytate content of the samples was determined following the method described by [11]. Four gram of the grinded sample was weighed into a beaker and was soaked in 100 ml of 2% HCl for 5 h and then filtered. 25ml of the filtrate was taken into a conical flask and 5 ml of 0.3% potassium thiocyanate solution was added. The mixture was titrated with a standard solution of Iron (III) chloride (FeCl_3) until a brownish yellow color persisted for 5 min. The concentration of the FeCl_3 was 1.04%w/v and Mole ratio of Fe (iron)to phytate = 1:1 100 x weight of sample concentrations of phytate phosphorous = Titre value x 0.064

Alkaloids

Alkaloids was estimated using the method by [18] Five grams of the samplehydrolysate was extracted in 200 ml of 10% acetic acid in ethanol. Samples was incubated for 4 h at room temperature, filtered before concentrating on a water bath. The extract was precipitated by the addition of drops of concentrated ammonium hydroxide and allowed to settle. The precipitate was washed with dilute ammonium hydroxide and then filtered. The residue which comprised of the alkaloid, was dried, and weighed. The alkaloid content was determined using the formula:

$$\text{alkaloid (\%)} = \text{final weight of sample/initial weight of extract} \times 100.$$

Bulk density

Five (5) gram of flour sample of the tiger nut was poured into a 100 ml measuring cylinder. The cylinder was tapped continuously until a constant volume was obtained. The bulk density (g/cm^3) was calculated as weight of flour (g) divided by volume of flour (cm^3). [19].

$$\text{Bulk density} = \frac{\text{weight of sample (g)}}{\text{volume of sample (ml)}}$$

Foam capacity (FC) and Foaming stability

FC was determined according to the method described by [20] using slurries that was prepared at 20, 40, or 60 mg/mL (protein weight basis) dispersions in 50 mL graduated centrifuge tubes containing 0.1 M phosphate buffer, pH 3.0, 5.0, 7.0, and 9.0. Sample slurry was homogenized at $20000 \times g$ for 1 min using a laboratory blender (Bl231japan). The capacity of the continuous phase to include air (FC) was determined as follows using the mean of measurements.

$$\text{Foam Capacity (FC)} = \frac{\text{Vol. after homogenization} - \text{Vol. before homogenization}}{\text{vol. before homogenization}}$$

Foam Stability (FS) The ability to retain air for a certain period (foam stability) was calculated by measuring the foam volume after storage at room temperature for 30 min and expressed as percentage of the original foam volume

$$\text{Foam stability (FS)} = \frac{\text{Volume after standing} - \text{Volume before whipping, ml}}{\text{vol. before whipping, ml}}$$

Water Absorption capacity (WAC)

The WAC was determined using the method of [20] with slight modifications. Protein sample (1 g) was dispersed in 10 mL distilled water in a 15 mL pre-weighed centrifuge tube. The dispersions were vortexed for 1 min, allowed to stand for 30 min and then centrifuged at $5000 \times g$ for 25 min at room temperature. The supernatant was decanted, excess water in the upper phase drained for 15 min and tube containing the protein residue was weighed again to determine amount of water retained per gram of sample.

Oil Absorption capacity (OAC)

The OAC was determined using the method of [11] with slight modifications. Protein sample (1 g) was dispersed in 10 mL pure canola oil in a 15 mL pre-weighed centrifuge tube. The dispersions were vortexed for 1 min, allowed to stand for 30 min and then centrifuged at $7000 \times g$ for 25 min at room temperature. The supernatant was decanted, excess oil in the upper phase drained for 15 min and tube containing the protein residue was weighed again to determine amount of water or oil retained per gram of sample.

Swelling index

Swelling index was determined according to the method by [21]. Two (2) grams of the flour sample was poured into a 50ml measuring cylinder and the volume it occupied was recorded. Already boiled water was added up to 50ml mark and the measuring cylinder was allowed to stand for 45mins after which the new volume of flour was recorded. The ratio of the initial volume to the final volume was taken as the swelling index.

$$\text{Swelling index} = \frac{\text{change in volume of sample (ml)}}{\text{Original volume of sample (ml)}}$$

Emulsifying properties.

Emulsifying activity index (EAI) and emulsion stability index (ESI) were determined using the method modified by [20]. Protein slurry of 10, 25, or 50 mg/ml each were mixed with 30 ml of deionized water. This protein solution was mixed with 10 ml of sunflower vegetable oil, and the pH was adjusted to 3, 5, 7, and 9. The mixture was homogenized at a speed of 1,000 rpm for 1 min. Fifty microlites (50 mL) of the aliquot of the emulsion were transferred (using pipette) from the bottom of the container at 0 and 10 min after homogenization and mixed with 5 ml of 0.1% sodium dodecyl sulphate (SDS) solution. The absorbance of the diluted solution was measured at 500 nm using spectrophotometer (UVIKON 930, BIO-TEK Kontron, Germany). This was calculated EAI and ESI using the method suggested by [11].

$$\text{Emulsifying activity index EAI} = \frac{2 \times 2.303 \times A_0}{0.23 \times \text{protein weight g}}$$

$$\text{Emulsifying stability index ESI (min)} = \frac{A_{10} \times Dt}{DA}$$

where A_0 is the absorbance at 0 min after homogenization; A_{10} is the absorbance at 10 min after homogenization; $Dt = 10$ min; and

$$DA = A_0 - A_{10}.$$

Least gelation concentration (LGC)

The LGC was determined according to the method of [20] with slight modification. 1 gram sample were suspended in water at different concentrations (2% to 20%, w/v, protein weight basis). The mixture was vortexed, placed in a water bath at 95 °C for 1 h, cooled rapidly under tap water and left in the refrigerator (4 °C) for 2-14 h. The sample concentration at which the gel did not slip when the tube was inverted was taken as the LGC.

Determination of protein Dispersibility

Dispersibility was determined by using the method described by [22]. 10g of the samples were weighted into a 100ml measuring cylinders and water added to make up to 100ml. The set- up was stirred vigorously and allowed to stand for three hours. The volumes of settled particles were taken and subtracted from 100.

$$\% \text{ Dispersibility} = 100 - \text{Volume of settled particle}$$

Protein solubility (PS)

The method by [32] was used and modified as follows. An aqueous solution (1% w/v) of protein sample was stirred for 30 min. With either 0.5 M HCl or 0.5 M NaOH, each solution was adjusted to the desired values (pH 3.0-8.0). The solution was centrifuged at 10000xg for 20 min. Modified Lowry method was used to determine the protein content using bovine serum albumin (BSA) as the standard. Percentage PS was expressed as: (protein content of each sample / total protein content) x 100. All determinations were carried out in triplicates.

In-vitro protein digestibility (IVPD) of Aduwa hydrolysates

In-vitro protein digestibility of the samples was measured according to the method described by Chavan *et al.* (2001). Two hundred and fifty milligrams of the sample were suspended in 15 mL of 0.1 M HCl containing 1.5 mg pepsin, followed by gentle shaking for 1 h at room temperature. The resultant suspension was neutralized with 0.5 M NaOH and treated with 4.0 mg pancreatin in 7.5 mL of phosphate buffer (0.2 M, pH 8.0). The mixture was shaken for 24 h at room temperature. The mixture was then filtered using Whatman No 1 filter paper and the residue washed with distilled water, air-dried, and used for protein determination using Lowry method [13] as described earlier.

Protein digestibility was obtained using the equation.

$$\text{In vitro protein digestibility (\%)} = \left(\frac{I - F}{I} \right) \times 100$$

where, I=protein content of sample before digestion

F=protein content of sample after digestion

Mineral analysis of Aduwa hydrolysates

The analyses for essential mineral elements were carried out by the atomic absorption spectrophotometric method described by [24] with modification. The sample (0.5g) was weighed into 75 ml digestion flask and 5 ml digestion mixture (10ml HNO₃ and 10ml HCl) added and digested at 150 °C until the solution becomes clear. It was cooled and 30 ml of distilled water added. The tube was vigorously stirred. The blank sample was prepared following the procedure describes earlier but with exception of the sample. A sample aliquot was then transferred to the Autoanalyzer (Technicon AAU model) for total mineral analysis at 420 nm. The left-over digest was used to determine the other elements (calcium, magnesium, and iron) on the Atomic Absorption Spectrophotometer (Perkin Elmer, model 402) while sodium and potassium were determined by flame photometry.

Data analysis

The result of three replicate experiments were analyzed for mean ± standard deviation. A one-way analysis of variance (ANOVA) and the least significance difference (LSD- turkey test) were carried out. Significance difference was accepted at P ≤ 0.05.

Results and discussion

Percentage yield of Aduwa protein hydrolysates.

The percentage oil recovery and material yield are shown in Table 1.

Sample	Quantity	Mean	Stdev	% Yield
APHpa (Pancreatin Hydrolysate yield)	Q1=200g Q2=92g, Q3=91.6g Yield= Q2/Q1 =0.46 Yield= Q3/Q1 =0.458	46.045.8	0.1	45.9
APHpe (Pepsin Hydrolysate yield)	Q1=200g Q2=75.2g, Q3=74.4g Yield= Q2/Q1 =0.376 Yield= Q3/Q1 =0.372	37.6037.2	0.2	37.4
APHpa+pe(Combined Hydrolysate yield)	Q1=500g Q2=222g, Q3=231g Yield= Q2/Q1 =0.444 Yield= Q3/Q1 =0.462	44.4046.2	0.9	45.1

Table 1. Percentage Yield of *Aduwa* protein hydrolysates

Key: APHpa= *Aduwa* protein Hydrolysates by pancreatin, APHpe= *Aduwa* protein hydrolysate by pepsin, APHpa+pe= *Aduwa* protein hydrolysate by pancreatin +pepsin combined

The percentage material yield of pancreatin hydrolysate APHpa (45.9%) and pepsin hydrolysate APHpe (37.4%) and APHpan+pe (45.1%) respectively showed that APHpa and APHpan+pe had better yield when compared to enzymatic pepsin hydrolysate samples APHpe. This observation could be due to peptide molecular sizes, peptide bonds that are being attacked and broken by enzymes during hydrolysis. The pancreatin hydrolysate had higher material yield and could be more economically viable to processors than pepsin enzymatic hydrolyzed peptides.

Proximate composition of *Aduwa* hydrolysates

Proximate composition of *Aduwa* hydrolysates is shown in Table 2.

Sample	Moisture Content (%)	Fat (%)	Crude Fiber (%)	Protein (%)	Ash content (%)	Carbohydrate (%)
APHpe	8.33 ^b ±0.208	0.14 ^c ±0.015	0.00 ^a ±0.00	79.31 ^b ±0.85	0.76 ^b ±0.02	11.46 ^b ±0.7
APHpan	8.91 ^a ±0.051	0.18 ^a ±0.012	0.00 ^a ±0.00	73.82 ^c ±0.11	0.64 ^c ±0.02	16.45 ^a ±0.14
APHpa+pe	8.96 ^a ±0.020	0.16 ^b ±0.006	0.00 ^a ±0.00	80.58 ^a ±0.86	0.85 ^a ±0.02	9.46 ^c ±0.84
LSD	0	0	0	0.418	0.62	0.06

Table 2. Proximate Composition of *Aduwa* protein hydrolysates

Mean value is from three determinations. Means followed by the same alphabetic on the column are not significantly different at $p > 0.05$. Key: isolate APHpa= *Aduwa* protein Hydrolysates by pancreatin, APHpe= *Aduwa* protein hydrolysate by pepsin, APHpa+pe= *Aduwa* protein hydrolysate by pancreatin +pepsin combined

Moisture content in food matrixes is one of the most important components of food processing and preservation. The moisture content is of direct economic importance to consumer, processor, and transporters. It is very significant; however, moisture affects the stability and quality of foods. The moisture content of the hydrolysate sample by all protease showed no significant difference. The moisture content of APHpe was (8.33%), APHpan (8.91%) and APHpan+pe (8.96%) respectively.

The protein content of *Aduwa* protein hydrolysate by pancreatin, pepsin and combined enzymes- pancreatin and pepsin significantly ($p > 0.05$) differ as sample protein hydrolysate are being made with different and combined enzymes. Crude protein content of *Aduwa* protein hydrolysate by combined enzymes APHpan+pe (80.50%) is significantly higher than separate enzyme hydrolysate, APHpan was (73.82%) and APHpe was (79.31%). The variation could be attributed to enzyme nature, activities, and possible cleaving site these proteases could have cleavage. peptide protein is an essential component of the diet required for the survival of both humans and animals. *Aduwa* protein hydrolysate can serve as a source of bio nutrient fortification. These could also serve as source of specific protein fractions for animal feed making at this hydrolysate state.

Fats are macronutrients, along with carbohydrates and protein. Fat is an important foodstuff for many forms of life and serves as both structural and metabolic functions. They are necessary part of the diet of both humans and animals and the most efficient form of energy storage. The crude fat content of the hydrolysate samples differ significantly. The pancreatin *Aduwa* hydrolysate has high fat content compared to APHpen (0.18%) and APHpan+pe (0.14%). [25], reported (9.63%) fat content on desert date kernel and this result was supported by [26]. This reported value by [25] is however high and far above hydrolyzed *Aduwa* samples. The low-fat content of the hydrolysates is an indication that it can be a good source material for food products required at low fat mix.

The ash content of the hydrolyzed sample analyzed differ significantly. *Aduwa* protein hydrolysate by combined enzymes APHpan+pe (0.85%) are significantly higher compared to APHpe (0.76%), and APHpan (0.64%) respectively. Since ash is the

index of mineral content, the combined hydrolysate meal has mineral contents or profile that could be physiologically important.

The Aduwa hydrolysates analyzed in this study contain no amount of crude fiber and differ significantly in all the samples ; APHpep (0.00%) , APHpan (0.00%) and APHpan +pe (0.00%) .The low values in crude fiber content could be because of the different proteases used on the concentrate samples .Low crude fiber content in nuts could lead to constipation if excess of it is being consumed as crude fiber enhances bowel movements Crude fiber is known to expand the inside walls of the colon, easing the passage of waste, and this makes it quite effective against constipation[27] .

Carbohydrates, alongside fats and proteins, are one of the three macronutrients in our diet with their main function being to provide energy to the body. Energy value of the Aduwa hydrolysates differs. Aduwa APHpan (16.45%) has higher energy values significantly compared to APHpan+pe (9.46%) and APHpep (11.46%). The low carbohydrate content in the hydrolysate sample might be due to the use of enzymes for hydrolysis, implying that Aduwa hydrolysate are not excellent source of carbohydrate rather peptides.

Mineral composition Aduwa hydrolysates

The mineral profile of protein materials from Aduwa is shown in Table 3.

Sample	Ca(mg/100g)	K (mg/100g)	Na (mg/100g)	Mg (mg/100g)	Fe (mg/100g)	Cu(mg/100g)	Zn(mg/100)
APHpe	0.64 ^a ±0.002	1.23 ^b ±0.002	1.18 ^b ±0.035	1.08 ^a ±0.035	0.20 ^b ±0.002	0.003 ^c ±0.00	0.12 ^c ±0.00
APHpan	0.85 ^b ±0.022	1.05 ^b ±0.004	2.06 ^c ±0.015	0.96 ^b ±0.015	0.15 ^b ±0.001	0.10 ^b ±0.004	0.24 ^b ±0.00
APHpa+pe	0.92 ^c ±0.055	1.29 ^c ±0.353	2.09 ^c ±0.163	1.09 ^c ±0.163	0.23 ^c ±0.125	0.09 ^b ±0.353	0.26 ^b ±0.20
LSD	0.101	0.32	0.101	0.101	0.023	0.32	0.003

Table 3. Mineral composition Aduwa hydrolysates

Mean values are readings from triplicate determinations; Means followed by the same alphabetic on the column are not significantly different at $p>0.05$

Key. AHPa= Aduwa protein Hydrolysates by pancreatin, AHpe= Aduwa protein hydrolysate by pepsin,AHpa+pe= Aduwa protein hydrolysate by pancreatin +pepsin combined

SAMPLE	Alkaloid mg/g	TPC mgCAT/g	Saponin mg/g	Tannin (mgCAT/g)	Phytate mg/g	Oxalate mg/100g
APHpa	Not detected	0.007 ^a ±0.00	0.012 ^a ±0.00	0.044 ^a ±0.00	Not detected	Not detected
APHpe	Not detected	0.055 ^b ±0.01	0.011 ^b ±0.00	0.022 ^b ±0.00	Not detected	Not detected
APHpa+pe	Not detected	0.084 ^c ±0.01	Not detected	0.009 ^c ±0.00	Not detected	Not detected
LSD		0.22	0.45	0.12		

Table 4. Anti -nutrients composition of Aduwa hydrolysates

Mean values are readings from triplicate determinations; Means followed by the same alphabetic on the column are not significantly different at $p>0.5$

Key.APM=AHPa= Aduwa protein Hydrolystaes by pancreatin, AHpe= Aduwa protein hydrolysate by pepsin,AHpa+pe= Aduwa protein hydrolysate by pancreatin +pepsin combined

Mineral calcium was significantly $p>0.05$ high in APH pa+pe (0.92mg/100g) APM compared to APHpa (0.85 mg/100g) and APHpe (0.64 mg/100g). Potassium content in APH pa+pe (0.92mg/100g) is significantly higher than APH pe (0.64 mg/100g) and APHpa (0.85 mg/100g). Similar trend was observed in mineral sodium, iron and zinc: Soduim had (APHpa2.06 mg/100g),

APHpe(1.18 mg/100g) and APH pa+pe(2.09 mg/100g), Iron had (APHpa (0.15 mg/100g), APH pe(0.20 mg/100g) and APH pa+pe (0.23 mg/100g), zinc had APHpa(0.24 mg/100g), APH pe(0.12 mg/100g) and APH pa+pe(0.26 mg/100g) ,manganese had (APHpa (0.96 mg/100g) , APH pe (1.08 mg/100g)and APH pa+pe (1.09 mg/100g) content respectively. Mineral copper in APHpa(0.105 mg/100g) was significantly different at ($p>0.05$) compared to APHpe(0.03 mg/100g) and APH pa+pe (0.09 mg/100g). The results show that Aduwa meal hydrolysate samples are rich in potassium, calcium, sodium, and magnesium, while other mineral such as copper which can helps the body form collagen, absorbs iron, plays a role in energy production and zinc plays a role in wound healing as well as treatment to diarrhea. The findings in this study agree with similar findings reported by[26] Supplementing these protein materials could curb child and adult Tetany osteomalacia and related diseases from due to lack of calcium. Potassium and sodium are electrolytes needed for the body to function normally and help in maintaining the fluid and blood volume of the body. Iron is a mineral that serves several important functions, its main function being to carry oxygen throughout our body and making red blood cells [28]

Anti Nutrients composition of Aduwa hydrolysates

Anti-nutrient composition of (*Balanites aegyptiaca del*) *aduw*a enzymatic hydrolysate are shown in Table 4. The presence of alkaloid disappeared and are absent in all enzymatic hydrolysate samples. These variations in the understudy might be due to the treatments employed. Alkaloid is an antimicrobial bio active characterized by bitterness 29(Ogori *et al* .2019) however, the alkaloid is reduced to zero in Aduwa hydrolysate samples. However toxic at a very high amount and may have physiological activities [25].

Total phenol content TPC are conjugated bioactive materials but varies depending on exposed sites [29]. The phenolic content in this study decreased significantly; APHpa (0.007mg GAE/g) APHpe (0.055 mg GAE/g) and APHpa+pe (0.084mg GAE/g) respectively. The results obtained in this study were lower than the values reported by [29] for soaked and roasted Aduwa samples. These indicates that enzymatic processing of *Aduwa* seed influenced phenolic profile content. The saponin contents of APHpa (0.002mg/g), APHpe (0.011 mg /g) and APHpa+pe (0.00mg/g) respectively under this study were significantly low and safe below lethal levels. Tannin content under this study reduced significantly as material samples were resolved enzymatically (APHpa (0.044 mgCAT/g), APHpe (0.022 mgCAT/g), APHpa+pe (0.009 mgCAT/g) .The phytate values under this study were completely absent APHpa(0.00mg/g), APHpe (0.00mg/g)and APHpa+pe(0.00mg/g). The phytate value obtained from Aduwa hydrolysates are lower than the lethal dose reported in other studies while the toxic effect of these anti-nutrients may not occur when these hydrolysates are consumed because their levels are not enough to elicit toxicity. Oxalate is another anti -nutrient moiety that causes intestinal. However, Oxalates were absent in all enzymatic hydrolysate samples. The values between 3-5mg/g have been pegged by [31] to be a lethal level. There was a significant decrease in hydrolysates, and these were within safety benchmark by[31]. Implying that enzymatic cleavage by pancreatin and pepsin had reducing effects on oxalate anti-nutritional factors

Functional properties of Aduwa (*Balanites aegyptiaca del*) seed meal, deffated meal, protein concentrate, isolate and hydrolysate

The Bulk density, WAC, OAC, LGC, and dispersibility of (*Balanites aegyptiaca.del*) *Aduwa* hydrolysates are shown in Table 5.

Sample	Bulk density(g/ml)	Swelling index (g/ml)	WAC(g/g)	OAC(g/g)	LGC (%)	Dispersibility (%)
APHpa	1.02 ^a ±0.15	3.45 ^a ±0.50	1.58 ^a ±0.06	1.72 ^b ±0.15	20.39 ^a ±0.6	62.12 ^b ±0.15
APHpe	0.12 ^b ±0.15	0.22 ^b ±0.21	1.72 ^a ±0.05	1.73 ^a ±0.50	14.36 ^b ±0.62	52.84 ^c ±0.14
APHpa+pe	0.11 ^c ±0.10	0.15 ^c ±0.10	0.17 ^c ±0.12	1.02 ^c ±0.02	14.32 ^b ±0.48	74.01 ^a ±0.42
LSD	0.000	0.0007	0.00017	0.0000	0.9000	1.1600

Table 5. Functional properties of Aduwa hydrolysates

Mean values are triplicate determinations: Means followed by the same alphabetic on the column are not significantly different at $p > 0.05$ Key.

APHpa= *Aduwa* protein Hydrolysates by pancreatin, APHpe= *Aduwa* protein hydrolysate by pepsin, APHpa+pe= *Aduwa* protein hydrolysate by pancreatin +pepsin combined

AHpan (1.02%) samples had significantly better packaging properties than APHpep (0.12%) and APHpan +pe (0.11g/ml), hence good weight and space relationship. The ability of biomaterial to absorb moisture and swell to a given capacity is influenced by hydrophilic or hydrophobic site exposure on their biomolecules. Swelling index from the Table 5 revealed that *Aduwa* APHpa 1.02 g/mL are significantly high than APHpe (0.12%) and APHpe+pa (0.11). This variation could be attributed to enzyme or the protease activities. The ability of protein material micelles to hold water molecules depends on the conformational position of the protein material, size, and shape. [32]. According to [33] this behavior is attributed to the hydrophilic and hydrophobic balance of the residual amino acid in the material. The WAC in APHpe (1.72 g/g) is significantly high than APHpan+pe (0.17g/g) and APHpa (1.58g/g). The variation observed could be attributed to the hydrolysis process. The OAC decreased ($p > 0.05$) from APHpa+pe(1.02)g/g to ,APHpa (1.72g/g) and then 1.73g/g in APHpe . The *Aduwa* hydrolysate by pancreatin and pepsin had the least OAC and are significantly different when compared to hydrolysate by combined enzymes APHpa+pe. However, this value did not agree with the value obtained from peas, chicken peas and lentils concentrate at these range (1.10-2.3g/g),[34] and walnut protein concentrate (2.50 g/g) [35]. This may suggest that *Aduwa* hydrolysate samples has good nonpolar amino acids, greater surface area of macro molecules, charges, and hydrophobicity properties. The LGC in APHpe (14.36) % and, APHpa (14.32) % are lower than APHpa+pe. The least gelation concentration of protein material confers gel formation through aggregation of denatured protein molecules. Gelation concentration helps in food system to ascertain degree of thickening and gelling especially in pudding and sources [36]. The ability of the hydrolysate samples to disperse easily in solution increased significantly at $p > 0.05$, APHpa+pe (74.01%) had the higher value when compared to in APHpe (52.84) % and, APHpa (62.12) %. The reconstitution ability of pepsin and pancreatin hydrolysate in aqueous medium were low compared to APHpa+pe. This observation maybe due to their bond sit resulting in their high percentage dispersion in water solution.

Protein solubility of *Aduwa* hydrolysate at different pH




Figure 1. Protein solubility of *Aduwa* protein isolate and hydrolysate at different pH

Key.APM= API= *Aduwa* proten isolate AHPa= *Aduwa* protein Hydrolysates by pancreatin , AHpe= *Aduwa* protein hydrolysate by pepsin,AHpa+pe= *Aduwa* protein hydroysate by pancreatin +pepsin combined

Figure 1 shows the solubility profile of *Aduwa* hydrolysate with respect to different pH (3, 5, 7 and 9) values. The results showed that the samples were most soluble at pH 4.0. The percentage soluble peptides decreased progressively as the pH value was adjusted from 4 to 8. The results showed that protein hydrolysate by combined enzyme, pancreatin and pepsin were least soluble at the very acidic pH value (pH 2-4). Ordinarily, the protein hydrolysates were expected to show better solubility at the acidic pH,

but the low solubility of the enzymatic Aduwa hydrolysates and at pH 2 when compared to the Aduwa protein meal may be attributed to high protein aggregation at the pH value, which reduced the solubility. Similar pattern of results was reported for okra seed meals and protein isolate [37]. Beyond pH 4.0, hydrolysate samples did show marked difference in the solubility, even as the pH value increased from 3-9. The protein hydrolysate had lowest protein solubility at pH 4.0 and thereafter increased progressively till pH 9.0, which agreed with the pattern of results reported for walnut protein [35]. The low values in solubility of the hydrolysates at pH 4.0 have helped to justify the iso-electric point of the hydrolysates. Usually, solubility decreases as the pH increases until it reaches the isoelectric point. The loss of electrostatic repulsive forces provides beneficial conditions for the formation of protein aggregates; high bulk density and large diameter of the aggregates results in precipitation of protein [38]. The difference between the pattern of solubility in protein hydrolysates may be due to the enzymatic hydrolysis. However, the low values and the pattern of solubility of the hydrolysate samples may be a disadvantage when considering its use as ingredients in acidic drinks.

Invitro protein digestibility of *Balanites aegyptiaca* del Aduwa protein hydrolysates

In-vitro protein digestibility of *Balanites aegyptiaca* del Aduwa hydrolysate is shown in Table 6.

Samples	Protein content before digestion	Protein content after digestion	<i>In-vitro</i> protein digestibility
	(%)	(%)	(%)
APHpan	54	7.43	84.93 ^b ± 0.10
APHpep	27	5.34	81.58 ^c ± 0.20
APHpan+pep	61	6.95	89.53 ^a ± 0.01
LSD			33.23

Table 6. Invitro protein digestibility of *Balanites aegyptiaca*. del Aduwa hydrolysates

Mean values are readings from triplicate determinations; Means followed by the same alphabetic on the column are not significantly different at $p > 0.05$

Key. AHPa= Aduwa protein Hydrolystaes by pancratin, AHpe= Aduwa protein hydrolysate by pepsin, AHpa+pe= Aduwa protein hydroysate by pancreatin +pepsin combined

The invitro protein digestibly increased significantly in APHpa+pe (89.53%) compared to APHpan (84.93%), and APHpe (81.58%). The high value of protein digestibility observed in hydrolysate samples may be due to their peptide fractions release, The Increase in *in vitro* protein digestibility experienced in hydrolysate samples may be due to the reduction in the levels of antinutritional factors.

Foaming capacity of Aduwa hydrolysates at different concentration and pH


Figures 1a, b and c show the influence of pH (3, 5, 7 and 9) and sample concentration (20, 40 and 60 mg/ml) on the foaming capacity of the samples. At the sample concentration of 20 mg/ml, the APHpe has high foaming capacity at pH3.0 and pH 5.0 respectively while the least foaming capacity was obtained at APHpa+pe and APHpan. The foaming capacity of Aduwa pancreatin hydrolysate APHpa and *Aduwa* pepsin hydrolysate APHpe decreased progressively as the pH of the solution increased from 3-9 at 20 mg/mL. The pattern was different in combined enzyme hydrolysate whereby the foaming capacity of the samples were relatively stable as the pH of the samples increased towards the basic region. The pattern of the results on enzymatic hydrolysate samples is in line with the increase in the net charge of the samples at the neutral and basic region, with the potentials to increase the net charge which eventually resulted in increase in protein-protein repulsion and a corresponding increase in the protein flexibility. When proteins become flexible, the tendency to accommodate more air bubbles increase and hence, an increase in the foaming capacity at the high pH values. Similar pattern of results was observed in the foaming capacities of fenugreek seeds,

bambara seed and walnut isolated proteins [35]. As the sample concentration was increased from 20 to 60 mg/ml, an increase in the foaming capacity of the APHpe was observed, basically in pH values 3, 7 and 9 but the foam formation at pH 5.0 remain substantially stable. For the APHpa and APHpe, an apparent increase in the foam capacity of the samples were observed up-to 60 mg/ml but decreased in values afterwards. A possible explanation for this pattern may that of protein crowding in APHpa result in from protein protein interactions. Although, an increase in the protein concentration is necessary to generate adequate foams; increase beyond 40 mg/ml may lead to generation of excess protein micelles that reduced the capacity to generate foams in the pancreatin and pepsin hydrolysate [39]


Foaming stability of Aduwa hydrolysates at different concentration and pH

Foaming stability is the ability of foam to keep its shape and volume over a specified period. This is very important because food material with good foaming stability could find applications in beverages, coffee, and baking industries. The foaming stability of *Aduwa* enzymatic hydrolysate with respect to variations in sample concentration (20, 40 and 60 mg/mL) and pH (3, 5, 7 and 9) values is shown in Figures 2 a,b and c.

A



B



C




Figure 2. a,b and c :Foaming capacity of aduwa hydrolysates at different concentration and pH

Means are readings from triplicate determinations. Means followed mean followed by the same alphabetic on the bars are not significantly different at $p > 0.05$

Key. AHPa= Aduwa protein Hydrolysates by pancreatin, AHpe= Aduwa protein hydrolysate by pepsin, AHPa+pe= Aduwa protein hydrolysate by pancreatin +pepsin combined

At 20 mg/mL sample concentration, the foam became increasing stable at the acidic pH (3 and 5) but increased progressively as the pH moved towards the basic region (7 and 9) for enzymatic hydrolysate samples. The results also revealed that values obtained for the foam stability were high in the APHpa+pe, APHpe and AHPa sample at 20 mg/mL sample concentration. This observation

or pattern in enzymatic hydrolysate samples may be attributed to the formation of stable molecular layers in the air-water interface that could have enhanced greater impartation of texture, stability, and more elasticity of foams. Similar pattern of results was reported for rapeseed by [40] . As the sample concentration increased from 20-60 mg/mL, the foam stability increased at the pH values 7 and 9, when compared with the acidic regions of 3 and 5 and this may suggest production of adequate charge densities at these pH values by the protein molecules which had made charges available to participate in the formation of strong interfacial membrane [41].At another observation, the result also showed that the foam stability was higher at sample concentration of 60 mg/ml, at a high pH values which also may suggest that the increase in the sample concentration is desirable in such that more protein molecules are produced to enhance the intermolecular cohesiveness of the foams formed [33]. The samples exhibited different pattern of foam stability with respect to the pH and varied sample concentration, which may be related to differences in the structural properties of the samples, especially the surface hydrophobicity in hydrolysate samples.

Emulsifying activity of aduwa hydrolysates at different concentration and pH.




Figure 3. a,b and c :Foaming Stability of aduwa hydrolysates at different concentration and pH.Means are readings from triplicates determinations. Means followed mean followed by the same alphabet on the bars are not significantly different at p>0.05

Key. AHPa= Aduwa Hydrolystaes by pancreatin AHpe= Aduwa hydrolysate by pepsin,AHpa+pe= Aduwa hydroysate by pancreatin +pepsin combined

Figures 3a, b and c show the emulsion activity of Aduwa protein enzymatic hydrolysate as functions of variations in pH (3,5 7 and 9) and concentrations of sample at 10, 15 and 50 mg/mL. At 10 mg/mL, the emulsion capacity of the APHpa, decreased from 3-9mL when compared to APHpe and APHpa+pe. The result also showed that the emulsion capacities of the enzymatic hydrolysate samples were low at high concentration of (25mg/mL and 50 mg/mL), respectively, this may be attributed to the

release of excess protein molecule which may have resulted in protein overcrowding or interaction and disruption in the interfacial properties [41].

Emulsifying stability of Aduwa protein meal, concentrate, isolates and hydrolysates at different concentration and pH

The potential of any protein to interact and bring together two immiscible phases such as oil and water and prevent phase coalescence is measured by emulsion stability [33]. Figures 4a, b and c




Figure 4 a-b: Emulsifying activity of Aduwa protein hydrolysates at different concentration and pH. Means are readings from duplicate determination. Means followed mean followed by the same alphabet on the bars are not significantly different at $p > 0.05$

Key.AHPa= Aduwa protein Hydrolysates by pancreatin, AHPe= Aduwa protein hydrolysate by pepsin,AHPa+pe= Aduwa protein hydrolysate by pancreatin +pepsin combined




Figure 5.a-b: Emulsifying stability of Aduwa protein meal, concentrate, isolates and hydrolysates at different concentration and pH. Means are readings from triplicate determinations. Means followed mean by the same alphabet on the bars are not significantly different at $p > 0.05$

Key.APM= AHPa= Aduwa protein Hydrolystaes by pancratin ,AHPe= Aduwa protein hydrolysate by pepsin,AHPa+pe= Aduwa protein hydroysate by pancreatin +pepsin combined

show the emulsion stability of enzymatic hydrolysate as function of varied pH (3,5,7,9) and sample concentrations 10, 25 and 50 mg/mL respectively. The emulsion stabilities of hydrolysate AHPa was high at 10 mg/ml, but the highest emulsion stability was obtained at pH 5 and pH9. But the emulsion formed at pH3, for AHPe was strong at 25 mg/mL sample concentration. However, at sample concentration of 50 mg/mL, pH 5.0 exhibited stronger AHPa+pe emulsion activities than at pH 3. The pattern of the emulsion stabilities in this study for the samples showed that sample concentration of 10 mg/mL and 50mg/mL are the threshold concentration for samples to create enough interfacial tensions to stabilize the emulsion formed by these hydrolysate samples.

Conclusion

The proximate, mineral, and phytochemical properties of the combined hydrolysate had better advantages over the pancreatin and pepsin hydrolysate samples. The solubility of enzymatically hydrolysate for both single and combined hydrolysate samples compounds from Aduwa at alkaline pH, showing average solubility score. WAC, OAC and LGC was improved only in the pepsin hydrolysate compound. The hydrolysate digest obtained pancreatin and pepsin show a significant difference relative to combined hydrolysate. Emulsifying properties were not improved, as proteases were combined. The crude protein, content IVPD and zero anti nutrient levels obtained from combined enzymatically hydrolysate digests shows that when included as ingredients in other

food products, they will improve nutritional quality, as they carry relevant amounts of peptides.

Conflict of Interest

The authors hereby declared no conflict of-interest

Acknowledgements

The author is grateful to the management of Federal university Gashua and Tet FUND for the support and the Department of Food Science and Technology, Federal University of Agriculture, Makurdi for the assistance during the study.

Reference

1. Nolsoe H, Undeland I (2009). The acid and alkaline solubilization process for the isolation of muscle proteins. *Food and Bioprocess Technology*, 2, 1–27
2. Ogori1 AF, Girgih AT, Eke MO, Abu O, Akinsola AF (2022) Phytochemical, anti-nutrient, invitro-protein digestibility and functional properties of (*Balanites Aegyptiaca* (L.) Delile) Aduwa protein meals, Protein Concentrate and Isolate Food Therapy and Health Care ,4: 10
3. Osemwota EC, Alashi AM, Aluko RE (2021) Comparative study of the structural and functional properties of membrane isolated and isoelectric pH precipitated green lentil seed protein isolates. *Membranes* 11: 694.
4. Girgih AT, Udenigwe CC, Aluko RE (2011) In vitro antioxidant properties of hemp seed (*Cannabis sativa* L.) protein hydrolysate fractions. *J Am Oil Chem Society* 88: 381–9.
5. Nancy DA, Aluko RE (2022) Acetylcholinesterase and butyrylcholinesterase inhibitory activities of antioxidant peptides obtained from enzymatic pea protein hydrolysates and theirultrafiltration peptide fractions. *J Food Biochem* 46: e14289.
6. Rosa CS (2000) Estudo das propriedades funcionais do colágeno obtido da pele de frango. Dissertaçãoem Mestrado em Ciência e Tecnologia de Alimentos. Universidade Federal de Santa Maria. Santa Maria: UFSM. Safari, R., Motamedzadegan, A, Ovi
7. Lempek TS, Martins VG, Prentice CH (2007) Rheology of surimi-based products fromfatty fish underutilized by the industry: Argentine croaker (*Umbrina canosai*). *Journal of Aquatic Food Product Technology* 16: 27-44
8. Obidah W, Nadro MS, Tiyafu GO, Wurochekke AU (2009) Toxicity of Crude *Balanites aegyptiaca* Seed Oil in Rats. *Journal of America Science* 5: 13-16.
9. Gbadamosi SO, Abiose SH, Aluko RE (2012). Solubilization amino acid composition and electrophoretic characterization of conophore nut proteins. *International food research journal* 19 :651-56
10. Aluko RE, Monu E (2003) Functional and bioactive properties of quinoa seed protein hydrolysates. *J. Food Sci* 68: 1254-58.
11. Reddy NR, Person MD (1994) Reduction in anti-nutrient and toxic compounds in plant food by fermentation. *Food research international* 27: 281-90
12. Lowry O H, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193: 265-75.
13. AOAC (2010) (Association of Official Analytical Chemists). *Official Methods of Analysis*.13th Edition of the Association of Official Analytical Chemists. USA, Washington D.C. Rockville, Maryland.
14. Ihekoronye PN, Ngoddy PO (1985) *Integrated Food Science and Technology for the Tropics*. Vol 1 Macmillian Publishes London. 236-44.
15. Peri, C., Pompei, C. (1971). Estimation of different phenolic groups in vegetable extracts. *Phytochemistry* 10: 2187-89.
16. Falade OS, Sowwnnie OR, Oladipo AT, Adewusi JK (2003) The level of organic acids in some Nigeria fruits and their effect on mineral availability in composite diet. *Pakistan journal of Nutrition* 2: 83-8.

17. Brunner JH (1984) Direct Spectrophotometer Determination of Saponin. *Journal of Analytical Chemistry*, 34: 1314-26.
18. Harborne JB (2005) *Phytochemical Methods*. Edit. London Chapman and Hall Ltd 49-188.
19. Oladele A. K and Aina J. O. (2007) Chemical composition and functional properties of flour produced from two varieties of tigernut (*Cyperus esculentus* African *Journal of Biotechnology* 6: 2473-76
20. Adebisi AP, Aluko RE (2011) Functional properties of protein fractions obtained from commercial yellow field pea (*Pisum sativum* L.) seed protein isolate. *Food Chemistry*, 128: 902-08.
21. Onwuka GI (2005) *Food analysis and Instrumentation theory and practice* 1st edit. Lagos. Naphthalin print PP3
22. Kulkarni KD, Kulkarni DN, Ingle UM (1991) Sorghum malt-based weaning food formulations: Preparation, functional properties, and nutritive value. *Food and Nutrition Bulletin*, 13 :1-7.
23. Tsou MJ, Kao FJ, Tseng CK, Chiang WD (2010). Enhancing the anti-adipogenic activity of soy protein by limited hydrolysis with Flavoenzyme and ultrafiltration. *Food Chemistry* 122: 243-48.
24. Amadou I, Amaza T, Young -Hui S, Geo -Wei I (2011) Chemical analysis and antioxidant properties of foxtail millet grain extract. *Songla. Journal of science and Technology* 33: 509-15.
25. Datti Y, Tijjani YA, Koki IB, Ali UL, Labaran M, et al. (2020). Phytochemical composition of desert date kernel (*Balanites aegyptiaca*) and the physical and chemical characteristics of its oil. *Biological and Pharmaceutical Sciences*, 11: 197-207.
26. Jock AA (2011) Physicochemical and Phytochemical Characterization of Seed Kernel Oil from Desert Date (*Balanites Aegyptica*), *Journal of Chemical Engineering and Bio-Analytical Chemistry*, 2: 49-61
27. Boakye AA, Wireko -Man FD, Agbonorhev JK, Oduro L (2015) Antioxidant activity, total plants, and phytochemical constituent in fruit. *International, Journal Food Research*, 221: 262-26.
28. Beard JL, Dawson HD(1997) Iron. In: O'Dell, B.L. and Sunde, R.A., Eds., *Handbook of Nutritionally Essential Minerals*, Marcel Dekker, New York 275-334.
29. Ogori AF, Girgih AT, Lukas H, Zhanibek Y, Anuarbek S, et al. (2019) Assessment of the phytochemical and functional properties of pre-treated aduwa (*balanites aegyptiaca*) seed meal flour, *Journal of Microbiol. Biotech. Food Science*, 9: 354-35
30. Sathe SK (1994) Solubilization and electrophoretic characterization of cashew secondary structures of soybean 7S and 11S globulins using AOT reverse antioxidant properties of pea seed (*pisum sativum* L.) enzymatic protein hydrolysate antioxidant fractions. *Journal of Agricultural and Food Chemistry*, 58: 4712-18.
31. FAO, (1985) World Health Organization. Energy and Protein requirements. Reports of joint FAO/WHO/UNU Expert Consultation. World Health Organization, Room.
32. Chavan U, McKenzie D, Shahidi F (2001) Functional properties of protein isolates from beach pea (*Lathyrus maritimus* L.). *Journal of Food Chemistry*.74: 177-87.
33. Malomo SA, He R, Aluko RE (2015) Structural and functional properties of hemp seed protein products. *Journal of Food Science*. 79: C1512-21.

34. Boye J, Zare F, Pletch A (2010) Pulse proteins: Processing, characterization, functional properties and applications in food and feed. *Food Research International*, 43: 414-31.
35. Mao X, Hua Y (2012) Composition, Structure and Functional Properties of Protein Concentrates and Isolates Produced from Walnut (*Juglans regia* L.) *Molecular Sciences*, 13: 1561-81.
36. Mohize MAK, Malomo S, He R, Aluko RE (2014) Structural and Functional Properties of Hemp Seed Protein Products. *J. Food Sci* 79: 1512-21.
37. Nnamezie AA, Akinsola AF, Gbadamosi SO (2021) Characterization of okra seed flours, protein concentrate, protein isolate and enzymatic hydrolysates *Food Production, Processing and Nutrition* 9: 48594864.
38. Singh RP, Anderson BA (2005) The major types of food spoilage: an overview. *Understanding and Measuring the Shelf-life of Food* 3-23.
39. Yu L, Haley S, Perret J, Hari M, Wilson J, et al. (2007) Free radicals scavenging properties of wheat extracts. *Journal of Agriculture and Food Chemistry* 50: 1619-24
40. Mahajan A, Dua S, Bhardwaj S (2002) Simple physical treatment as an effective radical scavenging peptide from fermented mussel sauce and its antioxidant properties. *Food Research International* 38: 175-82.
41. Ijarotimi OS, Malomo SA, Fagbemi TN, Osundahunsi OF, Aluko RE (2018) Structural and functional properties of *Buchholzia coriacea* seed flour and protein concentrate at different pH and protein concentrations. *Food Hydrocolloids* 74: 275-88.