

RESEARCH ARTICLE

Prevalence of Metabolic Syndrome among Urban Older Adults of Hyderabad Metropolitan City in South India

Sneha M¹, Arlappa N^{1*}, Suryanarayana P², Balakrishna N³, and Santosh VS²

¹Division of Community Studies, National Institute of Nutrition (ICMR) National Institute of Nutrition (ICMR), Hyderabad, India

²Department of Lipid Chemistry, National Institute of Nutrition (ICMR), Hyderabad, India

³Department of Bio-Statistics, National Institute of Nutrition (ICMR), Hyderabad, India

***Corresponding author:** Arlappa N, Scientist – E (Deputy Director), Division of Community Studies, National Institute of Nutrition (ICMR), Jamai-Osmania PO, Hyderabad – 500007, India, E-mail: arlappan@yahoo.com

Citation: Sneha M, Arlappa N, Suryanarayana P, Balakrishna N, Santosh VS (2017) Prevalence of Metabolic Syndrome among Urban Older Adults of Hyderabad Metropolitan City in South India. J Nutr Obes 1: 101

Article history: Received: 11 January 2017, Accepted: 02 February 2017, Published: 03 February 2017

Abstract

Background: Ageing is associated with decline in metabolism, physical activity and increased prevalence of non-communicable diseases like hypertension, diabetes and dyslipidemia. This study was carried out with the objective to assess the prevalence of metabolic syndrome (MS) among urban older adult population.

Methodology: A community based cross-sectional pilot study was carried out among 112 geriatric population (≥ 60 years and over) in Hyderabad metropolitan city by adopting random sampling procedure. Anthropometric measurements and blood pressure were recorded and lipid profile and fasting blood sugar were estimated. International Diabetes Federation (IDF) criterion was adopted for diagnosis of metabolic syndrome. Descriptive statistics like mean (SD) and prevalence were calculated using SPSS (Version 19.0). Chi-square test was performed to study the association between the prevalence of Metabolic Syndrome and different variables.

Results: The prevalence of hypertension, central obesity, and high blood sugar was 79.3%, 76.6%, 62.2% respectively. While the proportion of geriatric population with hypertriglyceridemia, hypercholesterolemia and low HDL was 24.3%, 10.8%, 23.4%, respectively. Similarly, the prevalence of metabolic syndrome among urban geriatric population was 54.1% and it was significantly associated with central obesity and high body mass index (BMI).

Conclusions: In general, the prevalence of metabolic syndrome was high among urban geriatric population. It may be attributed to rapid urbanization and high prevalence of non-communicable diseases. Therefore, adaptation of healthy lifestyles during early stages of adulthood is very essential for the prevention and control of metabolic syndrome and promotion of healthy ageing.

Keywords: Older Adults; Elderly; Metabolic Syndrome and Urban

Introduction

Most regions and countries in the world are experiencing unprecedentedly rapid demographic change [1]. As a result, world population increased more than double to 6.5 billion in 2005 [2] with an increase in the proportion of the elderly. This rapid increase in the proportion of elderly population could be attributed to considerable decline in mortality and fertility rates globally due to rapid advancements in health care and improved access to public health care facilities [3]. As fertility rates decline, the proportion of persons aged 60 and over is expected to double between 2000 and 2050 (11% to 22%), and their actual number will be more than triple, reaching 2 billion by 2050 [4]. The most rapid increases in older populations are currently happening in the developing world. Indian population has approximately tripled during the last 50 years, but the number of elderly Indians has increased more than fourfold. In 2001, the elderly population (60+) of India was 77 million. As per census 2011, the absolute number of elderly population has crossed the 100 million mark. This number is expected to increase to 323 million, constituting 20 per cent of the total India population by 2050 [5].

As people age, they are at an increased risk of disease and disability, and often, have increasing health needs. Ageing affects almost all the systems of the body, and is associated with several physiological, metabolic and psychological changes. This enhanced longevity is associated with burden of several chronic non-communicable diseases, such as cardiovascular diseases (CVDs), diabetes, osteoporosis, cataract, cancer etc., which lead to disability and increased burden not only on the health system but also on the economy. High prevalence of overweight, obesity and related co-morbidities are considered as major health problems

among older adults [6]. Ageing is an important and universal contributor to etiology of metabolic decline and related diseases which include type 2 diabetes, cardiovascular diseases, stroke [7] and Insulin resistance [8]. Abdominal obesity is commonly observed during ageing contributes to insulin resistance and metabolic syndrome. Medical concerns such as cardiovascular diseases, type 2 diabetes, metabolic syndrome risk factors and obesity have a significant direct relationship with low or non-exercise activity [9]. Metabolic Syndrome is associated with a 2-fold risk of CVD and a 5-fold risk of diabetes. Individuals suffering from Metabolic Syndrome have a 30%–40% probability of developing diabetes and/or CVD within 20 years, depending on the number of components present [10]. It was reported that the prevalence of MS has been increasing in the populations of the developing countries as a result of changes in lifestyles and dietary pattern [11,12]. According to the published data, the prevalence of MS among the Indian population was ranged from 9 to 48% [13-16].

The risk of development of metabolic syndrome is increased with advancing age, affecting about half of the adults aged ≥ 60 years [11]. Similarly, according to Modified NCEP ATPIII and IDF criterion, 47.1% (Men:40.7% & Women:54%) and 35.6% (Men:23.1% & Women:49%) of urban geriatric population of Kolkata had MS, respectively [17].

Data on Metabolic Syndrome among adult population are readily available in India. However, there is a paucity of such data among geriatric population in India, particularly among urban geriatric population of Hyderabad, a metropolitan city of South India. Therefore, keeping in view the magnitude of disease burden among the older population, a community based study was carried out during the year 2015 among the urban geriatric population of Hyderabad city with the objective of assessing the prevalence of metabolic syndrome and its determinants.

Methods

A community based cross-sectional pilot study adopting simple random sample was carried out among the urban geriatric population of Hyderabad metropolitan city during the year 2015. A total of 112 subjects of both genders were covered from the sampling frame of older adults (60 years and over) for the study. A pre-tested questionnaire was administered to elderly to obtain the data on lifestyle, physical activity, type of diet and history of non-communicable diseases.

Anthropometric measurements

Anthropometric measurements such as height, weight, waist circumference (WC) and hip circumference (HC) were recorded on all the selected older adults, using standard equipment and adopting standard procedure. Weight was measured with minimum clothing using SECA digital weighing scale (nearest to 0.1 kg). Height was measured using anthropometric rod, with the subject made to stand erect on a flat surface (without footwear) with feet together. The nutritional status of the older adults was assessed based on body mass index (BMI) classification, which is derived by weight in kilograms divided by height in meters' square [18]. The WHO recommended BMI cut-off values for Asian adults were used to calculate overweight and obesity [19], while the WC cut-off values of ≥ 90 cm and ≥ 80 cm were considered for men and women respectively, to calculate abdominal or central obesity. [20]

Clinical assessment

Blood pressure was measured thrice using an OMRON BP apparatus with the interval of at least 5 minutes between measurements. BP was recorded on the left arm of the subject in sitting position. Systolic and diastolic blood pressure was calculated as the mean of the three measurements. Fasting blood glucose and lipid profile was estimated from the subjects with 12 hours fasting.

Biochemical analysis

Fasting blood samples (6 ml) were collected from each subject by venipuncture for analysis of total plasma cholesterol, triglycerides (TG) and HDL. The plasma was separated from the whole blood by centrifugation and the lipid profile was estimated using kit methods (Bio-Systems). Fasting blood glucose was estimated by glucometer (Accu-Check Active Glucometer- Roche Diagnostics).

Definition of Metabolic Syndrome

Metabolic syndrome was defined as per the International Diabetes Federation (IDF) [21] criteria of presence of central obesity (Waist circumference for South Asian Men: ≥ 90 cm and South Asian Women: ≥ 80 cm) plus 2 of any of the following:

| Measure | Categorical Cut-off Points |
|--|---|
| Elevated waist circumference for Asian adults | Men: ≥ 90 cm Women: ≥ 80 cm |
| Elevated triglycerides (drug treatment for elevated triglycerides is an alternate indicator) | ≥ 150 mg/dL or 1.7 mmol/L or specific treatment for this lipid abnormality |
| Reduced HDL-C (drug treatment for reduced HDL-C is an alternate indicator) | Men: < 40 mg/dL or 1.03 mmol/L Women: < 50 mg/dL or 1.29 mmol/L |

| Measure | Categorical Cut-off Points |
|---|--|
| Elevated blood pressure (antihypertensive drug treatment in a patient with a history of hypertension is an alternate indicator) | Systolic BP \geq 130 or Diastolic BP \geq 85 mm Hg |
| Elevated fasting glucose (drug treatment of elevated glucose is an alternate indicator) | >100 mg/dL or 5.6mmol/L |

Source: Alberti *et al.* Circulation. 120(16):1640-5.0

Table 1: Criteria for Clinical Diagnosis of the Metabolic Syndrome

Ethical clearance and Consent

The study protocol was approved by the Scientific Advisory Committee of National Institute of Nutrition (NIN) and the ethical clearance was obtained from the Institutional Ethical Review Board. Written informed consent was obtained from the subjects after explaining them the objectives of the study.

Statistical analysis

Statistical analysis was performed using the Statistical Package for Social Science (SPSS) software (version 19.0) [22]. Descriptive statistics like mean (SD) anthropometric and bio-chemical parameters and the prevalence of non-communicable diseases among the elderly were calculated by age group. Comparison of mean anthropometric and bio-chemical variables across age groups was done using t-test. Chi-square test was performed to study the association between the prevalence of Metabolic Syndrome and different variables. P value of <0.05 was considered as statistically significant.

Results

The demographic and coverage particulars are presented in Table 2. A total of 112 (Men: 68.8% & Women: 31.3%) free living older adults were covered for the study. Most of the households were either nuclear (56.3%) or extended nuclear families (21%) and the proportion of the literacy among the elderly was about 94%. All the elderly were married and about 82% were living with their spouses. The previous occupation of the majority (59.5%) of the older adults was either government or private service and the source of livelihood among 51% of the elderly was their own income.

| Particulars | n | Percent |
|------------------------|-----|---------|
| Gender | | |
| Men | 77 | 68.8 |
| Women | 35 | 31.3 |
| Type of Family | | |
| Nuclear | 63 | 56.3 |
| Extended nuclear | 21 | 18.8 |
| Joint | 28 | 25.0 |
| Literacy status | | |
| Illiterate | 7 | 6.3 |
| Primary | 10 | 9.0 |
| Secondary | 10 | 9.0 |
| 10+2 | 36 | 32.4 |
| Graduate | 39 | 35.1 |
| Post graduate | 9 | 8.1 |
| Marital status | | |
| Married | 112 | 100.0 |
| Living with spouse | 92 | 82.1 |
| Living alone | 20 | 17.9 |
| Currently working | 8 | 7.1 |
| Past occupation | | |
| Laborers | 2 | 1.8 |
| Service | 66 | 59.5 |
| Business | 5 | 4.5 |
| Others (House wives) | 30 | 27.0 |

| Particulars | n | Percent |
|----------------------------------|----|---------|
| Main source of livelihood | | |
| Own income | 57 | 50.9 |
| Support from family | 39 | 34.8 |
| Support from govt/NGO | 5 | 4.5 |
| Others | 11 | 9.8 |
| Smokers | 22 | 19.4 |
| Non-smokers | 90 | 80.4 |
| Alcoholics | 28 | 25.0 |
| Non-alcoholics | 84 | 75.0 |
| Vegetarian | 55 | 49.1 |
| Non-vegetarian | 57 | 50.9 |
| Physical activity | 88 | 78.6 |
| No Physical activity | 12 | 21.4 |

Table 2: Coverage particulars of older adult population

About 22% of the older adults were smokers (both current and past) with the mean duration of smoking 19.4 years. Similarly, one fourth of the study subjects were alcoholics (both current and past) with the mean duration of consumption of alcohol 23.8 years. About 49% of the study subjects were vegetarians and a majority (78.6%) of the older adults were engaged in different physical activities such as walking, yoga and playing games (Shuttle badminton etc.).

Anthropometric measurements

The mean (SD) anthropometric measurements and indices of older adults by age groups are presented in Table 3. In general, all the mean anthropometric measurements and indices except for height and waist hip ratio (WHR) were lower in ≥ 70 years age group as compared to 60-69 years age group. The mean anthropometric measurements such as WC and HC were significantly ($p < 0.05$) higher among age group of 60-69 years as compared to ≥ 70 years age group.

| Particulars | Age group | n | Mean (SD) | p-value |
|---------------------------------------|-----------|-----|---------------|---------|
| Height (cms) | 60-69 | 73 | 159.0 (8.90) | 0.167 |
| | ≥ 70 | 38 | 162.65 (7.07) | |
| | Pooled | 111 | 160.2 (8.47) | |
| Weight (kgs) | 60-69 | 73 | 68.0 (9.91) | 0.847 |
| | ≥ 70 | 38 | 65.3 (10.61) | |
| | Pooled | 111 | 67.1(10.19) | |
| BMI [†] (kg/m ²) | 60-69 | 73 | 26.8 (3.32) | 0.669 |
| | ≥ 70 | 38 | 24.6 (3.67) | |
| | Pooled | 111 | 26.0 (3.59) | |
| WC [‡] (cms) | 60-69 | 73 | 93.6 (8.40) | 0.402 |
| | ≥ 70 | 38 | 90.4 (10.23) | |
| | Pooled | 111 | 92.5 (9.15) | |
| HC [†] (cms) | 60-69 | 73 | 99.2 (7.37) | 0.419 |
| | ≥ 70 | 38 | 92.7 (8.24) | |
| | Pooled | 111 | 97.0 (8.23) | |
| WHR [‡] | 60-69 | 73 | 0.95 (0.77) | 0.555 |
| | ≥ 70 | 38 | 0.98 (0.07) | |
| | Pooled | 111 | 0.96 (0.08) | |

BMI[†]: Body Mass Index; WC[‡]: Waist Circumference; HC[†]: Hip Circumference; WHR[‡]: Waist Hip Ratio

Table 3: Mean (\pm SD) of anthropometric parameters of urban older adults

Biochemical parameters and Blood Pressure

In general, all the mean (SD) biochemical parameters were higher among 60-69-year-old elderly as compared to elderly of ≥ 70 years. Similar trend was observed with respect to diastolic blood pressure (DBP), while, the mean systolic blood pressure (SBP) was higher among ≥ 70 -year-old elderly as compared to elderly of 60-69 years. However, no significant differences in mean biochemical parameters and blood pressure were observed between age groups (Table 4).

| Particulars | Age group | n | Mean | p-value |
|---------------------------|-----------|-----|----------------|---------|
| HDL (mg/dL) | 60-69 | 74 | 36.8 (9.84) | 0.029 |
| | ≥ 70 | 38 | 36.3 (7.24) | |
| | Pooled | 112 | 36.7 (9.00) | |
| Total Cholesterol (mg/dL) | 60-69 | 74 | 159.7 (37.31) | 0.785 |
| | ≥ 70 | 38 | 148.6 (38.05) | |
| | Pooled | 112 | 155.9 (37.77) | |
| Triglycerides (mg/dL) | 60-69 | 74 | 131.3 (71.63) | 0.159 |
| | ≥ 70 | 38 | 95.4 (52.54) | |
| | Pooled | 112 | 119.0 (67.67) | |
| Blood glucose (mg/dL) | 60-69 | 74 | 133.8 (49.84) | 0.357 |
| | ≥ 70 | 38 | 131.6 (32.43) | |
| | Pooled | 112 | 133.0 (44.51) | |
| SBP [†] mm Hg | 60-69 | 74 | 135.3 ((17.55) | 0.263 |
| | ≥ 70 | 38 | 137.8 (14.89) | |
| | Pooled | 112 | 136.2 (16.66) | |
| DBP* mm Hg | 60-69 | 74 | 83.1 (8.93) | 0.558 |
| | ≥ 70 | 38 | 82.0 (8.20) | |
| | Pooled | 112 | 82.7 (8.66) | |

SBP[†]: Systolic Blood Pressure; DBP*: Diastolic Blood Pressure

Table 4: Mean (±SD) bio-chemical parameters and blood pressure among the urban older adults

Components of metabolic syndrome

The prevalence of various non-communicable diseases among the urban older adults is presented in Table 5. In general, about 23% of the older adults had lower HDL levels. While about 11% and 24% elderly had higher cholesterol and triglycerides, respectively. Similarly, the proportion of the older adults with diabetes and hypertension was 62.2% and 79.3%, respectively. Likewise, majority of the older adults had central/visceral obesity (76.6%) and higher BMI (85.6%) values. The prevalence of overall obesity was significantly (p<0.05) higher among 60-69-year-old elderly as compared to elderly of ≥ 70 years. Similarly, the overall prevalence of metabolic syndrome among urban elderly was about 54%, and the prevalence was relatively higher in 60-69 years' age group as compared to older adults of ≥ 70 years.

| Particulars | Age group | n | percent | p-value |
|-------------------------------|-----------|-----|---------|---------|
| HDL (Low HDL) | 60-69 | 74 | 19.2 | 0.143 |
| | ≥ 70 | 38 | 31.6 | |
| | Pooled | 112 | 23.4 | |
| Total Cholesterol (≥200mg/dL) | 60-69 | 74 | 11.0 | 0.944 |
| | ≥ 70 | 38 | 10.5 | |
| | Pooled | 112 | 10.8 | |
| Triglycerides (≥ 150mg/dL) | 60-69 | 74 | 31.5 | 0.015 |
| | ≥ 70 | 38 | 10.5 | |
| | Pooled | 112 | 24.3 | |
| Blood glucose (≥ 100mg/dL) | 60-69 | 74 | 54.8 | 0.027 |
| | ≥ 70 | 38 | 76.3 | |
| | Pooled | 112 | 62.2 | |
| Hypertension | 60-69 | 74 | 79.5 | 0.950 |
| | ≥ 70 | 38 | 78.9 | |
| | Pooled | 112 | 79.3 | |
| Central Obesity | 60-69 | 74 | 87.7 | 0.000 |
| | ≥ 70 | 38 | 55.3 | |
| | Pooled | 112 | 76.6 | |
| BMI (≥ 23 kg/m ²) | 60-69 | 74 | 90.4 | 0.045 |
| | ≥ 70 | 38 | 76.3 | |
| | Pooled | 112 | 85.6 | |

| Particulars | Age group | n | percent | p-value |
|--------------------|-----------|-----|---------|---------|
| Metabolic Syndrome | 60-69 | 74 | 58.9 | 0.155 |
| | ≥ 70 | 38 | 44.7 | |
| | Pooled | 112 | 54.1 | |

Table 5: Prevalence (%) non-communicable diseases, the components of Metabolic Syndrome among the urban older adults

The association between the prevalence of Metabolic Syndrome and different variables is presented in Table 6. In general, the prevalence of MS was significantly ($p < 0.001$) associated with obesity, where the elderly with central obesity and overweight/obese had high prevalence of MS. However, no significant association was observed between the prevalence of MS and variables such as gender, age group, literacy status and lifestyles in this pilot study ($p > 0.05$).

| Variable | n | MS [†] | Chi-square | p-value |
|------------------------------|----|-----------------|------------|---------|
| Men | 77 | 53.2 | 0.147 | 0.701 |
| Women | 35 | 57.1 | | |
| Age group | | | 2.020 | 0.155 |
| 60-69 | 74 | 58.9 | | |
| ≥ 70 | 38 | 44.7 | | |
| Illiterates | 7 | 71.4 | 1.413 | 0.493 |
| Up to 10th Standard | 20 | 60.0 | | |
| College and above | 84 | 51.2 | | |
| Smokers | 22 | 59.1 | 0.236 | 0.627 |
| Non-smokers | 90 | 53.3 | | |
| Alcoholics | 28 | 50.0 | 0.300 | 0.584 |
| Non-alcoholics | 84 | 56.0 | | |
| BMI | | | 13.254 | 0.000 |
| 18.5 -23.0 kg/m ² | 16 | 12.5 | | |
| ≥ 23.0 kg/m ² | 96 | 61.5 | 40.500 | 0.000 |
| Waist Circumference | | | | |
| Normal | 26 | 0.0 | | |
| Central Obesity | 86 | 70.9 | 0.157 | 0.692 |
| Vegetarian | 55 | 56.4 | | |
| Non-vegetarian | 57 | 52.6 | | |
| Physical activity | 72 | 59.7 | 2.247 | 0.134 |
| No Physical activity | 40 | 45.0 | | |

[†]MS: Metabolic Syndrome

Table 6: The association between the prevalence of Metabolic Syndrome and different variables

Discussion

The current study on prevalence of metabolic syndrome among the free living urban geriatric population, perhaps, was carried out for the first time in Hyderabad metropolitan city. In general, the proportion of older adults of urban Hyderabad with MS was very high (54.1%), where a majority of them had central obesity (76.6%), hypertension (79.3%), diabetes (62.2%) and dyslipidemia according to the IDF criteria. Higher proportion of older adults in Hyderabad had MS as compared to their counterparts in Chennai (37.4% as per IDF), the other metropolitan city of South India while, as per the modified ATP-III and WHO criteria, the prevalence of MS among the older adults in Chennai was 45.9% and 45.5%, respectively [23]. Similarly, the corresponding figures of MS reported for the elderly in Tirupathi, another South Indian city was low (34.6%) [24]. Likewise, Prasad *et al* reported high prevalence of MS (40-65.6%) among urban dwelling elderly of Odissa, an Eastern Indian State [25]. While Sinha *et al* reported the prevalence of MS as 42.1% among the elderly residing in old age homes of Hyderabad [26].

As reported by the Tehran Lipid and Glucose Study (TLGS), the prevalence of MS among urban elderly in Iran was 41.9% [27] and the corresponding figures reported for the urban elderly in Iran and Taiwan was 55.4% [28] and 34.3% [29], respectively. Similarly, a Chinese study reported high prevalence of MS (46.3% to 62.9%) among rural elderly [30]. Hypertension, central obesity and diabetes mellitus are the major contributors for the risk of prevalence of MS in urban geriatric population of Hyderabad and similar contributors were reported for the prevalence of MS by other studies [24,29,31,32]. Thus, it is reiterated here that older

adults with any one of these risk factors should be cautious and adopt healthy lifestyles for the prevention of the increased risk of developing MS. In this present study, the prevalence of MS was higher in 60-69 years age group as compared to the elderly of ≥ 70 years and similar observations were made by Xiao et al. [30]. Similarly, the prevalence of MS was higher among the elderly women as compared to their men counterparts [24-30].

No significant association was reported between the prevalence of MS and the consumption of alcohol and tobacco in any form and the findings are consistent with findings of western studies [27,29,30]. Whereas, Sánchez-Villegas *et al* reported that drinking more than three alcoholic beverages per day was associated with high prevalence of MS [33]. Similarly, in the present study no significant association was observed between the MS and physical activity ($p > 0.05$). The probable reasons could be a majority of the elderly adopted practicing physical activity only after they were diagnosed with one or more non-communicable diseases and irregular or insufficient duration of physical activity.

In general, the prevalence of MS was reported to be higher among the urban elderly as compared to their rural counterparts, which could be attributed to the impact of urbanization. Urbanization significantly impacts the lifestyles through dietary and environmental influences with a consequent adverse effect on human health where the risk of development of lifestyle diseases is expected to be high [34-36].

Conclusion

The prevalence of MS is high among the urban geriatric population of Hyderabad, a South Indian city. Globally MS has taken up a greater dimension in urban geriatric health, where the prevalence of lifestyle diseases such as hypertension, diabetes mellitus, obesity etc. is high. Therefore, it is essential to take appropriate intervention measures like primordial prevention i.e. prevention of the development of risk factors for lifestyle diseases through health and nutrition education (HNE) and behavioral change communication (BCC). Similarly, the community is encouraged to adapt healthy lifestyles such as healthy eating, good sleep pattern, limiting smoking and alcoholism etc. during early stages of adulthood that is essential for prevention and control of metabolic syndrome and promotion of healthy ageing.

Limitations of the study

This study was carried out on pilot mode. Similarly, the information of dietary pattern among the older adults was not collected in the present study.

Acknowledgement

This work was supported by Department of Science and Technology (NO. SB/EMEQ-153/2013), Government of India. We are grateful to the Director of National Institute of Nutrition. We are also thankful to Prasad Undrajavarapu and Bhoja Raju Banavath for their support in completing the project.

References

1. Bongaarts J (2009) Human population growth and the demographic transition. *Philos Trans R Soc Lond B Biol Sci* 364: 2985-90.
2. United Nations (2007) World population prospects: the 2006 revision, New York, USA.
3. Suraiya I (1999) Assessing Nutritional Vulnerability in Older People in Developing Countries. *UN ACC-SCN News* 19: 18-21.
4. Ageing and Life-course (2014) World health Organization.
5. Population Census 2011 (2011) Census of India.
6. Rössner S (2001) Obesity in the elderly—a future matter of concern? *Obes Rev* 2: 183-88.
7. Ford ES, Giles WH, Dietz WH (2002) Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 287: 356-9.
8. Morley JE (2008) Diabetes and aging: epidemiologic overview. *Clin Geriatr Med* 24:395-405.
9. Hamilton MT, Hamilton DG, Zderic TW (2007) Role of Low Energy Expenditure and Sitting in Obesity, Metabolic Syndrome, Type 2 Diabetes, and Cardiovascular Disease. *Diabetes* 56: 2655-67.
10. Enas EA, Mohan V, Deepa M, Farooq S, Pazhoor S, et al. (2007) The metabolic syndrome and dyslipidemia among Asian Indians: a population with high rates of diabetes and premature coronary artery disease. *J Cardiometab Syndr* 2: 267-75.
11. Misra A, Khurana L (2008) Obesity and the metabolic syndrome in developing countries. *J Clin Endocrinol Metab* 93: S9-30.
12. Misra A, Vikram NK (2008) Factors, definitions, predictive value and Asian Indian ethnicity: Complexities of the metabolic syndrome. *Indian J Med Res* 127 : 293-6.
13. Misra A, Pandey RM, Devi JR, Sharma R, Vikram NK, et al. (2001) High prevalence of diabetes, obesity and dyslipidemia in urban slum population in northern India. *Int J Obes Relat Metab Disord* 25: 1722-9.
14. Gupta A, Gupta R, Sarna M, Rastogi S, Gupta VP, Kothari K (2003) Prevalence of diabetes, impaired fasting glucose and insulin resistance syndrome in an urban Indian population. *Diabetes Res Clin Pract* 61: 69-76.
15. Deepa M, Farooq S, Datta M, Deepa R, Mohan V (2007) Prevalence of metabolic syndrome using WHO, ATPIII and IDF definitions in Asian Indians: The Chennai Urban Rural Epidemiology Study (CURES-34) *Diabetes Metab Res Rev* 23: 127-34.
16. Sawant A, Mankeshwar R, Shah S, Raghavan R, Dhongde G et al. (2011) Prevalence of Metabolic Syndrome in Urban India. *Cholesterol* 2011: 1-7.

17. Sinha U, Mukhopadhyay B (2015) Metabolic syndrome in the urban elderly of Kolkata: effects of socio-demographic and lifestyle factors. *Asian J Gerontol Geriatr* 10: 1-6.
18. James WPT, Ferro Lyzzi A, Waterlow JC (1988) Definition of Chronic Energy Deficiency in Adults. *Eur J Clin Nutr* 42: 969-81.
19. World Health Organization. Regional Office for the Western Pacific (2000) The Asia-Pacific perspectives: redefining obesity and its treatment.
20. Alberti KG, Zimmet P, Shaw J, IDF Epidemiology Task Force Consensus Group (2005) The metabolic syndrome--a new worldwide definition. *Lancet* 366: 1059-62.
21. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JJ, et al. (2009) Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 120: 1640-5.
22. Statistical Package for the Social Sciences (2005) The SPSS guide to data analysis (Release 19.0). Chicago:USA.
23. Pradeepa R, Surendar J, Indulekha K, Chella S, Anjana RM et al. (2016) Prevalence of Metabolic Syndrome and its Association with Coronary Artery Disease among an Urban Elderly South Indian Population (CURES- 145). *J Assoc Physicians India* 64: 20-5.
24. Naik JL, Reddy KSN, Reddy BKC, Anwar Basha DA (2016) Prevalence and risk factors for metabolic syndrome in urban elderly: a community study from Tirupati, Chittoor district, Andhra Pradesh, India. *Int J Community Med Public Health* 3: 2106-111.
25. Prasad DS, Kabir Z, Das AK, Das BC (2012) Prevalence and risk factors for metabolic syndrome in Asian Indians: A community study from urban Eastern India *J Cardiovasc Dis Res* 3: 204-11.
26. Sinha N, Bhattacharya A, Deshmukh PR, Panja TK, Yasmin S, et al. (2016) Metabolic syndrome among elderly care-home residents in southern India: a cross-sectional study. *WHO South-East Asia J Public Health* 5: 62-9.
27. Hadaegh F, Zabetian A, Tohidi M, Ghasemi A, Sheikholeslami F, et al. (2009) Prevalence of metabolic syndrome by the Adult Treatment Panel III, International Diabetes Federation, and World Health Organization definitions and their association with coronary heart disease in an elderly Iranian population. *Ann Acad Med Singapore* 38: 142-9.
28. Frootan M, Mahdavi R, Moradi T, Mobasser M, Farrin N, et al. (2011) Prevalence of metabolic syndrome in an elderly population of Tabriz, Iran. *Endocrinol Metabol Syndrome S1*: 003.
29. Tsou MT, Chang BCC, Huang WH, Hsu CP (2014) Prevalence of Metabolic Syndrome and Risk Factor Analysis Among Urban Elderly in One Medical Center in Northern Taiwan. *Int J Gerontol* 8: 127-32.
30. Xiao J, Wu CL, Gao YX, Wang SL, Wang L, et al. (2016) Prevalence of metabolic syndrome and its risk factors among rural adults in Nantong, China. *Sci Rep* 6: 38089.
31. Arai H, Yamamoto A, Matsuzawa Y, Saito Y, Yamada N, et al. (2010) Prevalence of the metabolic syndrome in elderly and middle-aged Japanese. *J Clin Gerontol Geriatr* 1: 42-7.
32. Denys K, Cankurtaran M, Janssens W, Petrovic M (2009) Metabolic syndrome in the elderly: an overview of the evidence. *Acta Clin Belg* 64: 23-34.
33. Sánchez-Villegas A, Delgado-Rodríguez M, Alonso A, Schlatter J, Lahortiga F, et al (2009) Association of the Mediterranean dietary pattern with the incidence of depression: the Seguimiento Universidad de Navarra/University of Navarra follow-up (SUN) cohort. *Arch Gen Psychiatry* 66: 1090-8.
34. Adediran O, Akintunde AA, Edo AE, Opadijo OG, Araoye AM (2012) Impact of urbanization and gender on frequency of metabolic syndrome among native Abuja settlers in Nigeria. *J Cardiovasc Dis Res* 3: 191-6.
35. van der Sande MA, Milligan PJ, Nyan OA, Rowley JT, Banya WA, et al. (2000) Blood pressure patterns and cardiovascular risk factors in rural and urban Gambian communities. *J Hum Hypertens* 14: 489-96.
36. Ford ES, Giles WH, Dietz WH (2002) Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 287: 356-9.