

RESEARCH ARTICLE

Sustainable Fish Production Through Ecological and Fisheries Management of Canals in Indian Sundarbans (West Bengal)

Archana Sinha*, Tasso Tayung, Pranab Gogoi, Mitesh H. Ramteke, Aparna Roy, Arunava Mitra, Subhendu Mondal and Basanta Kumar Das

ICAR-Central Inland Fisheries Research Institute, Barrack pore, 700120, India

*Corresponding author: Archana Sinha, ICAR-Central Inland Fisheries Research Institute, Barrackpore, India 700120, Tel: 9433033950, E-mail: sinhaarchana@yahoo.com

Citation: Archana Sinha, Tasso Tayung, Pranab Gogoi, Mitesh H Ramteke, Aparna Roy, et al. (2021) Sustainable fish production through ecological and fisheries management of canals in Indian Sundarbans (West Bengal). J Aqua Sci Oceanography 1: 204

Abstract

The canal systems in Indian Sundarbans, West Bengal cover 907.33 ha area and offers a huge resources for fisheries development and livelihood. Naturally, fish enter to these canals from the source waters and form natural populations. Despite having vast resources, a little attention is paid to boost Inland fish production from canals. To understand the ecology and development of suitable practices/ techniques in canal systems, an effort was made for the sustainable fish production. Bharua canal a tidal fed canal connected with river Hooghly at Shibpur, Fraserganj of Sundarbans was selected to enhance fish production. The total length of canal is 2 km, average width of 45 m and water depth ranged from 3 to 8 feet. Three numbers of net partition system (NPS) were constructed with dimension of 50m × 45m covering an area of 2,250 m² each partition system. Water temperature (27.50±0.49°C), transparency (27.71±2.10 cm), Dissolved oxygen (5.61±0.13mgL⁻¹), pH (7.68±0.04), salinity (2.28±0.11gL⁻¹) were recorded during the trial. Diatoms (Bacillariophyceae) were dominated across the seasons. The quantitative abundance of phytoplankton and Zooplankton in the canal was 9522±2170 cellsl⁻¹ and 1918±97ind/l respectively. Rotifera shared dominance (45%) followed by crustacean nauplii (20%) and cladocera (12%). A total of 16 species of fish under 7 orders and 10 families were recorded. With the native fishes, fingerlings of Indian Major Carps *Labeo catla*, *Labeo rohita* and exotic carp *Ctenopharyngodon idella* were stocked @2 nos./m² in a ratio of 50: 40: 10 in a net partition system (NPS) installed in the canal. After a culture period of 180 days, the average size recorded for *Labeo catla*, *Labeo rohita* and *Ctenopharyngodon idella* were 265.61 ± 25.44 g, 184.63 ± 26.13 g and 221.57 ± 21.94g respectively. In addition to stocked fishes, considerable quantity of small indigenous fish (SIF'S) like *Puntius sophore*, *Puntius ticto*, *Amblypharngyodon mola*, *Salmostomo bacaila* were caught from NPS. In the present study, culture of fish in NPS showed feasible option in the canal. Hence, the NPS can be look upon as a potential culture practice to enhance the fish production and productivity from the unutilized canal resources. Adoption of this culture practice will provide a nutritional security, generate employment opportunity, and improve the livelihood and socioeconomic status of the rural populace of Indian Sundarbans.

Keywords: Canal Ecology; Carps Culture; Net Partition System; Indian Sundarbans

Introduction

Canals are the second most important (26%) source of irrigation covering 17 million ha in India (Agricultural Census 2010- 2011) which has considerable potential for fish production. The canal systems in Sundarbans covered area 907.33 ha and having vast potential for fish culture [1]. Various countries are producing fish from irrigations canals. Studies reported that around 16% of the freshwater fishery production comes from river Nile and its irrigation canals in Egypt while in Sudan, the fish biomass in minor canals of Gezira irrigation system with an average 660 kg/ha [2]. The practice of simple release of species *Oreochromis* spp., *Channa striata* and *Puntius gonionotus* in to irrigation canals reflected 350 kg/ha without supplementary feeding in Thailand [2]. Furthermore, the channelization or irrigation canals exhibit lower species diversity than the static water bodies which is primarily influenced by temperature and low primary producers [3, 4]. According to the published literature of FAO [5], fish biomass is 31% lower as compared to an unchannelised stream with 78% reduction of macro-invertebrate biomass. In Thailand, the extensive culture of bighead carp, grass carp and Nile tilapia in irrigation canals with the utilization of vegetable crops was successful [6, 2]. Physicochemical parameters in canals are strongly correlated with the phytoplankton abundance [7, 8]. The NO_2 , NO_3 , SiO_3 , HCO_3 , PO_4 , Ca^{2+} and Mg^{2+} are the key variables in the shaping of benthic fauna assemblage in Upper Ganga Canal (Matta, 2015); zooplankton primarily dominated by rotifers, cladocerans and protozoans in Ganga canal [10]. Limnological variations of Sirhind feeder channel (Hanumangarh) characterized by shallow with turbid, alkaline and well oxygenated water [10]. The study on fish culture practices in Indian canals are limited. In order to understand the ecology and development of suitable techniques in canal systems; research effort is necessary to achieve the sustainable production from these resources. Fisheries from canals are poorly documented. The information on trophic status of canal habitat, biotic community structure and hydrological parameters are lacking. Hence, the present study aimed to generate baseline data of canal ecosystem and development of suitable enhancement practices in canal ecosystem of Sundarbans, West Bengal.

Materials and Methods

The current study was carried out in Bharua canal which is located in Shibpur, Fraserganj of Sundarbans ($21^{\circ}36'39.3''\text{N}$ $88^{\circ}15'21.6''\text{E}$) West Bengal India, and is a tidal fed canal connected with river Hooghly. The total length of canal is 2.0 km and width are ~ 45 m, and having a water depth ranges from 3 to 8 feet. Three sampling locations were chosen at an equal distance with the station 1 at near river mouth (S1), then in the amidst of the canal (S2) and station 3 at the end point of the canal from river mouth (Figure 1). Sampling was carried out at bimonthly from April 2018 to March 2019, constituting three different seasons [11] i. e. pre-monsoon, monsoon and post-monsoon.




Figure 1: A Map of Bharua Canal, Bakkhali

Environmental Parameters Measurement

For physical and chemical analysis of water, with the help of standard water sampler based on the design of 'Ruttner water sampler [1], water samples were collected from subsurface (0.5 m depth) and instantly shifted into clean polythene bottles (1.0 L) from the predefined locations. Water temperature ($^{\circ}\text{C}$), transparency (cm), pH, dissolved oxygen (mgL^{-1}), specific conductivity (m S/cm), salinity (ppt), total alkalinity (mgL^{-1}), and total hardness (mgL^{-1}) were measured *in-situ*. For the analysis of other parameters such as total dissolved solids (TDS), and nutrients (phosphate, nitrate, sulphate, silicate), water samples were brought to the laboratory in cold conditions. A degree centigrade (-10 to 50 $^{\circ}\text{C}$) thermometer (P-601466) were used for measuring temperature, digital pH meter (Hanna Instruments) for recording water pH, a digital conductivity meter (Multiline P4-82362) for measuring specific conductivity, and Secchi disk was used for recording transparency (Strickland and Parsons, 1972). Analysis of total dissolved solids, dissolved oxygen, total alkalinity, total hardness, salinity and nutrients were determined by following the standard methods APHA (2012).

Sediment Quality Parameter

For sediment analysis, sediment samples were collected at about 30 cm depth and transferred in polythene bags, and brought to the laboratory for further analysis. For the chemical analysis, the samples were initially dried, made into finely powdered and sieved. The particle size of sediment was determined by hydrometer method. Value of sediment pH was determined by glass electrode pH meter in water using sediment/water ratio of 1:10. Specific conductivity calculated by conductivity meter. Organic carbon in the sample was determined by using wet oxidation method of Walkley and Black [12]. The available phosphorous and available nitrogen in the sample was determined by Brays II method [13] and alkaline potassium permanganate method [14], respectively. And, free calcium carbonate was measured volumetrically with HCL soluble CaCO_3 method (Ref.).

Phytoplankton Sample Collection

Phytoplankton samples were collected by filtering 50 L sub-surface water from the selected locations using plankton net (20 μm silk bolting mesh). The collected samples were preserved by adding 4% buffered formalin solution to it. For the quantitative analysis of plankton, a Sedgwick Rafter counting cell was used. The total number of phytoplankton cells present was calculated and expressed in terms of cells per litre (cells l^{-1}) [15]. Phytoplankton was identified by employing a trinocular microscope (Nikon Eclipse-50i) and with the help of published standard literatures [16, 17]. The genera of plankton were arranged according to the AlgaeBase, [18].

Fish diversity

Fish specimens were collected by experimental fishing using gears such as cast net, seine net, gill net and bamboo traps in the canal. In addition, fish samples were also collected from the local fishermen and markets nearby in the canal. The collected fish samples were preserved in 10% formalin solution. Identification of fishes was done by standard taxonomic keys and literatures [19, 20, 21].


Livelihoods and Ecosystem services

Data was also collected regarding the contribution of canal fisheries in the livelihoods of the rural populace living adjoin to the Bharua canal. A semi-structured interview schedule was prepared and data was collected through direct questioning. Data was collected from 20 households and also from the community leaders of that area. Ecosystem services rendered by the canal was also listed out based on the Millennium Ecosystem Assessment, 2005 synthesis report.

Installation of Net Partition System (NPS)

In collaboration with the local stakeholders of the Bharua canal, a net screen barrier as a net partition system (NPS) was installed in the canal. Three net partitions were constructed with dimension of $50\text{m} \times 45\text{m}$ (L \times B); each covering an area of $2,250 \text{ m}^2$. Net partition system is a kind of pen structure made from locally available bamboo poles and HDPE net. The erected structure of net

partition system is shown in Figure 1. Fingerlings of Indian major carps *Labeo catla*, *Labeo rohita* and exotic carp *Ctenopharyngodon idella* with average weight of 5.83 ± 0.13 g, 5.65 ± 0.33 g and 6.50 ± 0.41 g were stocked @2nos./m² with the stocking ratio of 50: 40: 10 respectively. The fishes were provided with an artificial feed (28% crude protein and 5% crude lipid) twice a day @4% of body weight for 180 days.



BAC: Bacillariophyceae; Cya: Cyanophyceae; Chl: Chlorophyceae; Cos: Coscinodiscophyceae; Eug: Euglenophyceae; Tre: Trebouxiophyceae; Zyg: Zygnematophyceae; Xan: Xanthophyceae; Din: Dinophyceae; Med: Mediophyceae; Syn: Synurophyceae

Figure 1(a): Compositions of phytoplankton in Bharua canal, Sundarbans

Growth performance analysis was carried out at bimonthly interval. The growth performances as recorded by means of length and weight of individual fish to determine weight gain (WG), relative growth rate (RGR %), specific growth rate (SGR %), and survival rate (SR %). These parameters were calculated as follows: $WG (g) = W_f - W_i$; $RGR (\%) = (W_f - W_i) / W_i \times 100$; $SGR (\%/day) = (\ln W_f - \ln W_i) / t \times 100$; Where, W_i and W_f are initial and final weights (g), respectively, and t is the time of the experiment (days); survival rate (%) = (no. of fish harvested / no. stocked) \times 100. Total fish yield was calculated as yield (kg) = Avg. final weight \times total no. of survivors.

Statistical Analysis

Parameters in this study were analysed using SPSS (version 23). Data are shown as means \pm standard deviation. Fish diversity has been measured by the number of species (species richness) and by using the three indices - (1) Shannon–Weiner index (H') (Shannon and Weiner, 1949): $H' = -\sum [P_i \times \log (P_i)]$; Where 'P_i' is the proportion of the individuals belonging to the 'ith' species and H' is the species diversity (2) Margalef richness index (d) (Margalef, 1958): $d = (S - 1) / \log N$, where N = total number of individuals and S = total number of species and (3) Pielou's evenness or equitability index (J') (Pielou, 1977): $J' = H' / \log (S)$; where H' = Shannon–Weiner diversity and S = total number of species. Similar diversity indices were also followed for phytoplankton population. One way analysis of variance (ANOVA) was performed to understand the significant heterogeneity of phytoplankton between seasons.

Results and Discussion

Environmental parameters measurement

Physical and chemical parameters of water

Seasonal variations in the physical and chemical parameters of Bharua canal are presented in Table 1. Salinity variation was wide in canal between the season's ranges from 0.37 – 3.54 ppt and highest salinity was recorded during pre-monsoon season. The pH of surface water ranged from 7.27 – 8.5. Overall, the pH of the canal was alkaline. Dissolved oxygen ranges from 4.4 – 6.5 mgL⁻¹, were favourable for good production of fish. Total alkalinity in Bharua canal ranges from 64 – 111 mgL⁻¹. So over all, the canal had a very good "buffering" capacity.

Table 1: Physico-chemical parameters of Bharua canal, Sundarbans (W.B)

Parameters	Pre-monsoon	Monsoon	Post-monsoon
Depth (cm)	121.67 ± 6.01	175.00 ± 6.45	310.00 ± 15.68
Air temperature (°C)	26.10 ± 0.06	29.40 ± 0.14	26.08 ± 0.05
Water temperature (°C)	26.70 ± 1.35	28.45 ± 0.06	27.35 ± 0.06
Transparency (cm)	27.70 ± 1.88	40.00 ± 1.87	15.43 ± 2.56
Specific conductivity (m S/cm)	6.08 ± 0.16	5.74 ± 0.17	0.93 ± 0.01
TDS (mgL ⁻¹)	536.67 ± 50.44	5003.50 ± 417.91	657.90 ± 44.39
pH	7.75 ± 0.05	7.95 ± 0.03	7.34 ± 0.03
DO (mgL ⁻¹)	5.70 ± 0.12	6.28 ± 0.11	4.85 ± 0.17
Free CO ₂	7.07 ± 0.23	5.35 ± 0.39	8.60 ± 1.15
Total Alkalinity (mgL ⁻¹)	130.00 ± 2.89	66.00 ± 1.35	108.25 ± 1.55
Salinity (ppt)	3.29 ± 0.16	3.18 ± 0.16	0.39 ± 0.01
Total hardness (mgL ⁻¹)	164.67 ± 4.33	705.25 ± 18.28	138.50 ± 13.84
Total N (mgL ⁻¹)	0.08 ± 0.01	0.56 ± 0.06	0.04 ± 0.00
Nitrate (mgL ⁻¹)	0.01 ± 0.00	0.01 ± 0.00	0.02 ± 0.00
Total phosphate (mgL ⁻¹)	1.90 ± 0.55	0.49 ± 0.03	2.57 ± 0.86
Phosphate - P (mgL ⁻¹)	0.07 ± 0.01	0.15 ± 0.01	0.06 ± 0.00
Silicate (mgL ⁻¹)	5.41 ± 0.35	7.07 ± 0.15	4.38 ± 0.77
Sulphate (mgL ⁻¹)	2.98 ± 0.41	145.21 ± 2.67	0.57 ± 0.16

The values are means ± SD

Soil Quality Parameters

Sediment size fraction of Bharua canal showed that major percentage of the sediment is contributed by silt (43.50 ± 1.76%) followed by sand (42.50 ± 0.87%) and clay (14.00 ± 2.61%). Organic carbon content in soil was low to medium (1.58 ± 0.07%); which means there is a less organic matter accumulation at the bottom. The soil pH of the canal was 7.08 ± 0.025. In this canal, an accumulation of available nitrogen and phosphorus were found in good quantity, especially available phosphorus. Sediment quality parameters of Bharua canal is presented in Table 2.

Table 2: Sediment quality of Bharua canal, Sundarbans

Parameters	Value
Sand (%)	42.50 ± 0.87
Silt (%)	43.50 ± 1.76
Clay (%)	14.00 ± 2.61
pH	7.08 ± 0.025
Specific conductivity (mS/cm)	7.35 ± 0.29
Available nitrogen (mg 100g ⁻¹)	18.62 ± 0.74
Available P ₂ O ₅ (mg 100g ⁻¹)	3.23 ± 0.47
Organic carbon (%)	1.58 ± 0.07
Free CaCO ₃ (%)	1.44 ± 0.16

The values are means ± SD

Plankton Diversity of Canal

The Bherua canal exhibited significant heterogeneity of plankton abundance across the seasons ($p < 0.05$). A total of 71 species (63 genera) of phytoplankton were recorded. Out of 11 algal groups, Bacillariophyceae (33%) excelled as major microfloral community. Algal group Cyanophyceae (28%) emerged as second conspicuous group (Figure 1). On the whole, three algal groups Bacillariophyceae, Cyanophyceae, and Chlorophyceae showed major contributions to the total phytoplankton community. The diatoms were dominated in terms of diversity (31 species) dominated by pennales. Seasonal abundance of phytoplankton revealed peak during post-monsoon season ($10,110 \pm 2696$) and lowest in pre-monsoon season (6385 ± 1233). Eight groups of zooplankton were recorded where Rotifera was dominant (45%) followed by Crustacean nauplii (20%) and Cladocera (12%). (Figure 2). The abundance of zooplankton observed maximum in monsoon season ($1,089 \pm 646$). The quantitative spectrum of phytoplankton and Zooplankton was $9,522 \pm 2170$ cells $^{-1}$ and $1,043 \pm 410$ ind. $^{-1}$, respectively in the canal. The Shannon-Weiner diversity (H') of phytoplankton was maximum in monsoon (3.2 ± 0.23) and lowest during pre-monsoon (3.1 ± 0.11). While, H' and Margalef richness (d') of zooplankton showed maximum during post-monsoon season. The richness and Shannon diversity indicated a rich phytoplankton diversity with a value of > 3.1 in the canal. Pielou's evenness index (J') reflected an even distribution of phytoplankton with mean value 0.88.

Fish Diversity of Bharua Canal

A total of 17 fish species recorded under 8 orders and 11 families and is presented on table 3. The catch structure revealed the dominance of family Cyprinidae (59%) followed by Bagridae (13%), Ambassidae (9%), Channidae (7%), Notopteridae (4%), Mastacembelidae (2%), Gobiidae (2%), Megalopidae (1%), Schilbeidae (1%), Anabantidae (1%) and Megalopidae (1%) respectively. The major component of fish catch was contributed by small indigenous fish (SIF'S) like *Puntius sophore*, *Puntius ticto*, *Amblypharngyodon mola*, *Rasbora daniconius* and *Salmotomo bacaila*. Seasonal fish diversity was observed highest during monsoon season. The fish diversity indices of Bharua canal is presented in figure 2.




Figure 2: Compositions of zooplankton in Bherua canal

(Diversity- Tasso)

Contribution of The Canal On Livelihoods and for Rendering Ecosystem Services

Bharua canal is located in Shibpur village in Namkhana Block of Sundarbans. Scattered human settlement is found by the side of the canal. The people living by the side of the canal often harvest fishes from the canals by using various traps and gears. People put bamboo made rectangular shaped traps like *Guni*, *Chero*, *Atal*, *Chokhia* or Triangular shaped trap *Banki/Mugri* to harvest fishes from marshy areas, water channels or canals in Sundarbans [22]. It was reported by the sampled populace that during monsoon, they are getting better catch and more types of high value fishes which lead an average income of INR 3850/- during monsoon months (June, July, August, September). However, other than the monsoon months, during lunar periods (*Bhara Kotal*) the people are getting more fish from this canals which contributes around INR 1260/- in their monthly income (during October-April). A similar study conducted in Frezarganj area of Sundarbans shows that fish catch from the water channels, canals, water bodies, paddy fields contribute 15.65 percent in the household income of the fishermen [23]. It was also reported by the sampled population that; the women of the adjoining households often collect Small Indigenous Fishes (SIFs) or prawns through small dragnets/scoop nets for the household consumptions.

Canals are rendering important ecosystem services both directly and indirectly. Fisher et al. 2009 has defined ecosystem services 'the aspects of ecosystems utilized (actively or passively) to produce human well-being'. The ecosystem services rendered by the canal has given below:

1. Provisioning services:
 - a. Source of fresh water
 - b. Provide 'Fin fish' and 'shell fish' as food
 - c. Irrigation water for crop production
 - d. Aquatic weeds as leafy vegetables
 - e. Agricultural diversity
 - f. Drainage of excess water
2. Regulating services:
 - a. Flood control
 - b. Carbon sequestration
 - c. Habitat for rare aquatic weeds and animals
 - d. Protection for coastal storm surges
 - e. Aquifer recharge/ ground water recharge
3. Cultural services
 - a. Site for ecological or environmental education
 - b. Human relaxation, walking trails
4. Supporting services
 - a. Primary productivity
 - b. Hydric soil development

Table 3: Fish species diversity of Bharua canals of Indian Sundarbans, along with their IUCN status and utilization

Sl. No.	Species	Order	Family	ICUN status	Utilization
1	<i>Channa punctata</i> (Bloch, 1793)	Channiformes	Channidae	LC	FF & OF
2	<i>Channa striatus</i> (Bloch, 1793)			LC	FF & OF
3	<i>Pethia ticto</i> (Hamilton, 1822)	Cypriniformes	Cyprinidae	LC	OF
4	<i>Puntius sophore</i> (Hamilton, 1822)			LC	OF
5	<i>Rasbora daniconius</i> (Hamilton, 1822)			LC	OF
6	<i>Amblypharyngodon mola</i> (Hamilton, 1822)			LC	FF
7	<i>Salmostoma bacaila</i> (Hamilton, 1822)			LC	FF
8	<i>Chanos chanos</i> (Forsskal, 1775)	Gonorynchiformes	Chanidae	NE	FF
9	<i>Macrogathus pancalus</i> (Hamilton, 1822)	Synbranchiformes	Mastacembelidae	LC	FF & OF
10	<i>Parambassis ranga</i> (Hamilton, 1822)	Perciformes	Ambassidae	LC	FF & OF
11	<i>Anabus testudineus</i> (Bloch, 1792)		Anabantidae	DD	FF
12	<i>Glossobogius giuris</i> (Hamilton, 1822)		Gobiidae	LC	FF & OF
13	<i>Mystus gulio</i> (Hamilton, 1822)	Siluriformes	Bagridae	LC	FF
14	<i>Mystus vittatus</i> (Bloch, 1794)			LC	FF & OF
15	<i>Pachypterus atherinoides</i> (Bloch, 1794)		Schilbeidae	LC	
16	<i>Notopterus notopterus</i> (Pallas, 1769)	Osteoglossiformes	Notopteridae	LC	FF & OF
17	<i>Megalops cyprinoides</i> (Broussonet, 1782)	Elopiformes	Megalopidae	DD	FF

Where, LC = Least count; NE = Not evaluated; DD = Data deficient

FF = Food fish; OF = Ornamental fish

Growth performance of fishes stocked in net partition system

The detail growth performance of stocked fishes is given table 4. The intent of present study was to observe the feasibility of Indian major carps and exotic carp culture in the slightly saline water canal of Sundarbans. After 180 days of culture, the avg. mean weight of *L. catla*, *L. rohita* and *C. idella* was 265.61 ± 25.44 g, 184.63 ± 26.13 g and 221.57 ± 21.94 g respectively. *L. catla* performed well in canal system than the other two species and attended highest weight gain (Fig. 2). Relative growth rate (RGR %) were highest in *L. catla* followed by *C. idella* and *L. rohita*. Specific growth rate (SGR %) of *L. catla* was 2.10 ± 0.33 followed by *C. idella* 1.95 ± 0.06 and *L. rohita* 1.91 ± 0.06 respectively. Highest survival % were recorded in *C. idella* 89.7 ± 2.02 % followed by *L. rohita* 82.85 ± 2.71 whereas, lowest in *L. catla* 77.53 ± 1.60 .

Table 4: Growth performance of fishes stocked in Net Partition System

Parameters	<i>Labeo catla</i>	<i>Labeo rohita</i>	<i>Ctenopharyngodon idella</i>
Initial weight (g)	5.83 ± 0.13	5.65 ± 0.33	6.50 ± 0.41
Final weight (g)	454.85 ± 46.35	380.73 ± 19.55	339.88 ± 29.17
Weight gain (g)	449.02 ± 46.38	375.08 ± 19.67	333.38 ± 28.98
Relative Growth Rate (RGR %)	7765.89 ± 871.77	6975.69 ± 832.85	5205.32 ± 400.70
Specific Growth Rate (SGR %)	2.40 ± 0.06	2.34 ± 0.05	2.19 ± 0.05
Survival rate (%)	77.53 ± 1.60	82.85 ± 2.71	89.7 ± 2.02

The values are means ± standard errors (n = 3)

Previous studies were conducted on seed raising of IMC as stocking material for wetlands. The fingerlings of *L. rohita*, *Cirrhinus mrigala* and *Hypophthalmichthys molitrix* reared up to the advance fingerling size for 59 days in pen with 4% body weight of supplementary feeding. The fishes attended the size from 50 to 71.5 g, 60 to 74 g and 30 to 75 g, respectively [24]. The advanced fry of IMC i.e. *L. rohita* and *Labeo catla* were reared in pen at Sareni wetland of Uttar Pradesh for 120 days. The mean weight attended by the fishes was 2.88±1.41 g to 57.2±13.84 g and 2.1±1.16 g to 67.38±25.79 g respectively [25].

In the present study, the fish yield in net partition system was recorded around 2484.34 kg from three partition unit, it means approximately 3,680 kg of fish biomass per hectare (Table 5). In Bihar, the study was conducted in ox-bow lakes to produce table size fish of IMC from pens by stocking advanced fingerlings, the result showed a yield of 3362-3962 kg/ha within a six month of culture period [26]. The fish yield in the present study found to be similar with their report. However, fish yield in the net partition system showed much higher than the yield (2106 and 1780 kg/ha-1) reported from pen culture in two Wetlands of Assam [21].

Table 5: Yield of the Stocked fishes in Net Partition System in Canal of Sundarbans

Species	No. of fishes stocked	No. of fishes Harvested	Yield (kg)	Yield(kg/m ²)
<i>Labeo catla</i>	6750	5233	1390.01	0.206
<i>Labeo rohita</i>	5400	4474	826.02	0.122
<i>Ctenopharyngodon idella</i>	1350	1211	268.31	0.040
Total	13500	10918	2484.34	0.368

The economic analysis of net partition system is presented in table 6. In the present experiment, benefit-cost ratio for the Net Partition System operation for rearing single crop was estimated to be 1.16, which is in agreement with the studies conducted in the wetland of Assam by [21] and flood plain wetland of Uttar Pradesh [25]. The return on investment estimated was 0.16.

Literature on culture of Indian major carps (IMC) in pen from Indian waters is limited whereas there is no report about pen culture in canal system. Previous studies were conducted for the culture of IMC seed for raising stocking material in the wetlands. [27] reported that *in-situ* raising of IMC fingerlings in pen enclosures in the floodplain wetlands of Uttar Pradesh is economically feasible. [25] reported that intervention through pen culture resulted in the increase in fish productivity of Sareni Jheel of Uttar Pradesh from 310 to 833 kg ha⁻¹. Inland fish production can sustainably increase by rational stocking in inland open water bodies such as wetlands, reservoirs and canals [28, 29, 30].

Result of the present study showed that culture of Indian Major Carps and exotic carp in net partition system in the canals of Sundarbans may provide the suitable alternative to the tribal fishers of Sundarbans to improve their livelihood and socioeconomic status.

Table 6: Economics of IMC & EC rearing in Net Partition System in Bharua canal

	Particulars	Rate (₹)	Unit	Amount (₹)
	Capital costs			
	Cost of bamboo poles (number)	300	45	13500
	Cost of HDPE net screen (meter)	400	50	20000
A	Sub total			33500
B	Capital costs per crop (assuming that the materials will last for 3 crops)			11166.67
	Operational costs			
	Cost of fingerlings (kg)	200	27	5400
	Cost of Feed (kg)	32.5	2000	65000
	Labour charges for construction of NPS	350	7	2450
C	Sub-total			72850
	Total costs (B+C)			84016.67
	Gross income			
	Sale value/ price of fish			
	<i>L. catla</i> (kg)	120	463.34	55600.80
	<i>L. rohita</i> (kg)	120	275.34	33040.80
	<i>C. idella</i> (kg)	100	89.44	8944
D	Sub-total			97585.60
E	Net income (Gross income - Total costs)			13568.93
	Benefit-cost ratio			1.16
	Return on investment			0.16

Conclusion

As the canals system in Sundarbans are holds great potential for the fish production but since long time they are unutilized or underutilized. The canal is a source of livelihood for the local populace and also mitigate their nutritional needs. Moreover, the present study showed that culture of fish in net partition system in canal is a feasible option. Adoption of net partition system in canal system can provides the alternate livelihood option to the fisher's community as the Indian major carp has good demand in local markets in addition to the native fishes of the canal.

References

1. Das Sarkar S, Naskar M, Gogoi P, Raman RK, Manna RK, et al. (2019). Impact assessment of barge trafficking on phytoplankton abundance and Chl a concentration, in River Ganga, India. *Plos One* 14: e0221451
2. Anon (2003) Fisheries in irrigation systems of Arid Asia, *FAO Fisheries Technical Paper No. 430*. Rome. 150 p.
3. Tarplee WHJnr, Louder DE, Weber AJ (1971) Evaluation of the effects of channelization on fish populations in North Carolina's coastal plain streams. North Carolina Wildlife Resources Commission, Internal Report 13 p.
4. Daget, J (1976) La production de poisson de consommation dans les ecosystemes irrigues. In: Arid land irrigation in developing countries-environmental problems and effects. Edited by E.B. Worthington, Pergamon Press, 295-301.
5. Anon (1990) Fish production in irrigation canals – A Review. *FAO Fisheries Technical Paper No. 317*. Rome. 111 p.
6. Little D, Muir J (1987) A guide to integrated warm-water aquaculture, Institute of Aquaculture, University of Stirling, Stirling, Scotland 238 p.
7. Agnieszka, Napiorkowska-Krzebietke (2014) Phytoplankton of artificial ecosystems-an attempt to assess water quality. *Archives of Polish Fisheries* 22: 81-96.
8. Bishnoi RK, Sharma BK, Durve VS, Sharma LL (2013) Primary productivity in relation to planktonic biodiversity in a stretch of Ganag Canal (Rajasthan). *Universal J Environ Res and Technol* 3: 266-72.
9. Bishnoi RK, Sharma BK (2016) Planktonic variations in lotic water body of Shri Gangasagar District, (Rajasthan). *Int J Fauna and Biol Stud* 3: 134-9.
10. Sharma M, Srivastava D (2016) Phytoplankton diversity and its ecology in Sadul Branch Sirhind feeder canal (Hanumangarh, Rajasthan). *Int J Pure & Appl Biosci* 4: 168-71.
11. Chaudhuri K, Manna S, Sarma SK, Naskar P, Bhattacharyya S, et al. (2012) Physicochemical factors controlling water column metabolism in Sundarban estuary, India. *Aquatic Bio systems* 8: 26.
12. Jackson ML (1973) Soil chemical analysis. New Delhi, India. Prentice Hall.
13. Baruah TC, Borthakur HP (1997) A Textbook of Soil Analysis. Vikas Publishing House Pvt. Ltd., New Delhi.
14. Subbiah BV, Asija GL (1956) A rapid procedure for the determination of available nitrogen in soil, *Cur Sci* 25: 259-60.
15. Santhanam P, Velayutham R, Jegathesan G (1989) A manual of freshwater ecology. Daya Publishing House, New Delhi
16. Belinger GE, Sigee CD (2010) Freshwater algae: Identification and use as 801 bioindicators. Blackwell publishing, p. 271.
17. Prescott GW (1962) Algae of the Western Great Lakes area with an illustrated key to the genera of desmid and freshwater diatoms. Revised Edn. W.M. C. Brown Company Publishers. Dubuque, Iowa. p. 10 0 0.
18. Guiry MD, Guiry GM (2019) AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>. Accessed 6 Aug 2019

19. Talwar PK, Jhingran AG (1991) Inland fishes of India and adjacent countries. Vol-1 and Vol-2. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, Bombay and Calcutta, pp 1063
20. Jayaram KC (1999) The Freshwater Fishes of the Indian Region. Narendra Publishing House, New Delhi, pp 551
21. Gorai BK, Sugunan VV, Jha BC (2006) Raising of Stocking Materials of Indian Major Carps in Pen Enclosures in Selected Floodplain Wetlands of Assam, India. *Asian Fisheries Sci* 19: 185-97.
22. Roy A, Sinha A, Manna RK, Aftabuddin MD, Das SK (2020) Traditional knowledge of the fishermen community of Indian Sundarbans: An assessment of rationality and effectiveness. *Indian J Fish* 67: 94-101.
23. Roy RK, Manna A, Sinha SK Das, S Majumder, DK Biswas, et al. (2016) A roadmap to ensure food and nutritional security in coastal Sundarbans. B. Mondal, D. Sarkar, S.D. Mukhopadhyay, S. Ghosh, B. Chowdhury, S.C. Roy (Eds.), *Family Farming Challenges and Opportunities*, Renu Publishers, New Delhi 218-28
24. Bhaumik U, Nath D, Saha S, Mitra A, Bandhopadhyay S (200) Demonstration of pen culture technology. In: Annual report 1999-2000. Central Inland Fisheries Research Institute, Barrackpore, India, 57-58 p.
25. Alam A, Joshi KD, Das SCS, Jha DN, Srivastava K, et al. (2017) Enhancing fish productivity through pen culture: a case study in Sareni wetland of Uttar Pradesh. *Indian J Fisheries* 64: 8-13.
26. Rai SP, Sigh RC (1990) Potential of pen culture in Ox-bow lakes. In: Jhingran, A. G., Unnithan, V.K., & Amitabha, G. (Ed.) *Proceedings of the Contribution to the fisheries of Inland open water systems in India*. Inland Fisheries Society of India Publication, Barrackpore, India. 13 p.
27. Das SCS, Alam A, Jha DN, Kumar V, Srivastava K, et al. (2017) Raising of stocking materials in pen enclosure in a floodplain wetland of Uttar Pradesh. *J Inland Fisheries Society of India* 49: 13-9.
28. Beveridge, Malcome CM (1984) Cage and pen farming-Carrying capacity models and environmental impact. *FAO Fisheries Technical Paper* 255, Rome. 131p.
29. Kutty MN, Campbell D (1987) Pen culture (Enclosure culture) as an aquaculture system. UNDP-FAO, Nigerian institutes of oceanography and marine research project RAF/82/009, Nigeria. 43p.
30. Baluyut EA (1989) Aquaculture systems and practices: A selected review. UNDP-FAO Report (ADCR/Rep/89/43), Rome. 1-80 p.
31. APHA (2012) In: Rice EW, Baird RB, Eaton AD, Clesceri LS (eds) *Standard methods for the examination of water and wastewater*, 22nd edn. American Public Health Association (APHA), American Water Works Association (AWWA) and Water Environment Federation (WEF), Washington, DC, USA
32. Fisher B, Turner RK, Morling P (2009) Defining and classifying ecosystem services for decision making. *Ecological economics* 68: 643-53.
33. Froese R, Pauly D (2021) FishBase. www.fishbase.org. Electronic version accessed 01/05/2021. Margalef, D.R., 1958. Information theory in ecology. *General Systems* 3: 36-71.

-
34. Matta G (2015) Effect of water quality on phytoplankton ecology of Upper Ganga Canal. *Int J Sci and Eng Res* 6: 762-8.
 35. Millennium Ecosystem Assessment (2005) *Millennium Ecosystem Assessment Synthesis Report*. Island Press, Washington, DC.
 36. Pielou EC (1977) *Mathematical ecology*. Wiley, New York, p 385.
 37. Shannon CE, Weiner W (1949) *The mathematical theory of communication*. University of Illinois, Illinois, p 125
 38. Simpson EH (1949) Measurement of diversity. *Nature* 163: 688.
 39. Strickland JDH, Parsons TR (1972) *A practical handbook of seawater analysis*. Bulletin, vol 167. Fisheries Research Board of Canada, Ottawa, p 326