

Unilateral Agenesis of the Superior Trapezius Muscle: A Case Report and Review of Literature

Marre' Brunenghi G^{1*} and Ruzzini L²

¹U.O. Orthopedics and Traumatology Giannina Gaslini Institute 5, Largo G. Gaslini 16147 Genova (Ge) Italy

²U.O. Orthopedics Bambino Gesù Palidoro Pediatric Hospital Via Torre di Palidoro, snc 00050 Fiumicino (Rome) Italy.

***Corresponding author:** Marre' Brunenghi, U.O. Orthopedics and Traumatology Giannina Gaslini Institute 5, Largo G. Gaslini 16147 Genova (Ge) Italy, Tel: +393357593927, E-mail: giorgiomarre@gaslini.org

Citation: Marre' Brunenghi G, Ruzzini L (2021) Unilateral Agenesis of the Superior Trapezius Muscle: A Case Report and Review of Literature. J Pediatr Dis Neonatal Care 2: 105

Abstract

We report the clinical features and radiological findings of a patient, who presented with unilateral partial agenesis of upper right trapezius, without any limitation of normal daily life. Since the majority of other reported cases of isolated partial or total trapezius agenesis are cadaveric occasional findings, we can assume that management is not required.

Keywords: Trapezius Muscle; Agenesis; Aplasia; Clinical Observation

Introduction

Trapezius is the largest of the dorsal trunk muscles and arises from the medial third of the superior nuchal line, the external occipital protuberance, the nuchal ligament and C7-T12 spinous processes [5]. It can be divided into three parts giving to its insertions: the superior or descending part that inserts onto the lateral third of the clavicle; the middle or transverse part that inserts onto the acromion and the inferior or ascending part that inserts on the spine of the scapula [1].

These three parts have also different functions such as the elevation and rotation of the scapula (superior part), the retraction of the scapula (middle part) and the depression and rotation of the scapula (inferior part) [3].

This muscle is innervated by the accessory nerve and by the C2, C3 and C4 ventral roots nerve fibers of the cervical plexus. Vascular supply is usually by the dorsal scapular artery or less commonly from the second part of the subclavian artery [7]. Contributions come also from the posterior intercostal arteries, from the occipital artery and thyrocervical trunk [1].

Reports of muscle absences are extremely rare. The most frequently involved muscle is pectoralis major with or without associated defects of pectoralis minor [1]. Congenital absence of upper limb muscles are mostly reported in association with pectoralis major defects as a component of Poland's Syndrome [2].

Trapezius muscle agenesis in association with other upper limb malformations is a rare event, and the isolated aplasia of trapezius either partial or complete is even more uncommon [3].

In literature 10 cases of trapezius muscle aplasia have been reported and only 6 of these were isolated upper limb anomalies [1, 3-11]. Of this cases just 1 was a clinical report [11]. The others were occasional findings in cadaveric cases [1, 3, 5, 7, 10].

We report the clinical features and radiological findings of a patient who presented with unilateral partial agenesis of upper right trapezius.

Case Report

A 9-year-old boy consulted us, complaining shoulder asymmetry. He did not report other problems; he was active and conducted a normal life. No family history of significance and no history of shoulder traumas were reported.

Physical examination revealed absence of the right superior trapezius (Figure 1). The remaining shoulder-girdle muscles were normal. A full range of movement was present in the cervical spine and in both shoulders. Forward flexion of the right shoulder was full but weaker than the left shoulder. External and internal rotation were full and normal in power. The right scapula was less stable than the left one and the "scapularis' pendulum" (the movement of the scapula during elevation and lowering of the upper limb) was asymmetric. There was no other physical abnormality on clinical examination.

Radiographs and Computed tomography (CT) of the right shoulder showed normal findings (Figure 2).

MRI of the right shoulder confirmed agenesis of the superior trapezius (Figure 3).



Figure 1: Clinical aspect

Physical examination revealed absence of the right superior trapezius




Figure 2: CT images

Radiographs and Computed tomography (CT) of the right shoulder showed normal findings



Figure 3: MR images

MRI of the right shoulder confirmed agnesis of the superior trapezius

Discussion

Muscle aplasia or hypoplasia are very rare conditions that usually affects pectoralis major and/or pectoralis minor in the Poland's Syndrome [12]. The etiology of this Syndrome is accredited to be a temporary blood supply interruption of the subclavian artery during the first gestational period [13]. In literature there are few reports of some variants of the Poland's Syndrome including also trapezius agnesis [6, 8, 9]. The theory at the base of this inclusion is that the trapezius muscle has its vascularisation from a branch of the subclavian artery [14].

Rarely [15] agnesis of trapezius could be associated to agnesis of sternocleidomastoid muscle.

Familial cases of agnesis of trapezius muscle are described [8, 16, 17] but our patient had not any relatives with similar problem.

Isolated absence of trapezius muscle is an extremely uncommon situation.

The biological basis for the agnesis of trapezius muscle is not well established; it remains a matter for speculation: genetic causes, congenital aberrations, embryological anomalies, anomalous blood supply, accessory nerve injury and atrophy of a muscular nervous centre can be all implicated in the aetiology of trapezius aplasia or hypoplasia [1, 3, 7].

Our case represents a rare manifestation of isolated, monolateral, partial trapezius aplasia. We are not aware of the causes underlying superior trapezius agnesis in our patient, even though it has been speculated that both local interruption in muscle blood supply and developmental anomalies can be the aetiology of isolated muscle aplasia [3].

The only concern of our patient is the shoulder asymmetry. As in our case, in fact, superior trapezius aplasia does not affect 10-year-old boy daily activities [11] and also sport activity in adult patients [17]. So we can speculate that, if isolated, this anatomical variation does not interfere with daily life and management is not required.

In conclusion, patients presenting with shoulder asymmetry, or shoulder impaired function should have trapezius agenesis added to the list of differential diagnoses.

Acknowledgements

We have to thank Dr. Paolo Violino (Radiologist of the Cuneo's Hospital) for the Computed Tomography and the Magnetic Resonance's scans.

References

1. Garbelotti SA Jr, Rodrigues CF, Sgrott EA, Prates JC (2001) Unilateral absence of the thoracic part of the trapezius muscle. *Surg Radiol Anat* 23: 131-3.
2. Castilla EE, Paz JE, Orioli IM (1979) Pectoralis major muscle defect and Poland complex. *Am J Med Genet* 4: 263-9.
3. Nooij LS, Oostra R (2006) Trapezius Aplasia: Indications for a Dual Developmental Origin of the Trapezius Muscle. *Clin Anat* 547-9.
4. Adams SB Jr, Flynn JM, Hosalkar HS, Hunter J, Finkel R (2003) Torticollis in an infant caused by hereditary muscle aplasia. *Am J Orthop* 556-8.
5. Allouh MZ, Mohamed A, Mhanni A (2004) Complete unilateral absence of trapezius muscle. *McGill J Med* 8: 31-3.
6. Debeer P, Brys P, De Smet L, Fryns JP (2002) Unilateral absence of the trapezius and pectoralis major muscle: A variant of Poland syndrome. *Genet Couns* 13: 449-53.
7. Emsley JG, Davis MD (2001) Partial unilateral absence of the trapezius muscle in a human cadaver. *Clin Anat* 14: 383-6.
8. Gross-Kieselstein E, Shalev RS (1987) Familial absence of the trapezius muscle with associated shoulder girdle abnormalities. *Clin Genet* 32: 145-7.
9. Horan FT, Bonafede RP (1977) Bilateral absence of the trapezius and sternal head of the pectoralis major muscles. A case report. *J Bone Joint Surg* 59A: 133.
10. Rahman HA, Yamadori T (1994) An anomalous cleido-occipitalis muscle. *Acta Anat (Basel)* 150: 156-8.
11. Witbreuk MM, Lambert SM, Eastwood DM (2007) Unilateral hypoplasia of the trapezius muscle in a 10-year-old-boy: a case report. *J Pediatr Orthop B* 16: 229-32.
12. Urschel HC Jr (2009) Poland syndrome. *Semin Thorac Cardiovasc Surg* 21: 89-94.
13. Bavinck JN, Weaver DD (1986) Subclavian artery supply disruption sequence: hypothesis of a vascular etiology for Poland, Klippel-Feil, and Mobius anomalies. *Am J Med Genet* 23: 903-18.
14. Nurettin Yiyit, Turgut Isitmangil, Coskun Ozturker (2014) The abnormalities of trapezius muscle might be a component of Poland's syndrome. *Medical Hypotheses* 83: 533-6.
15. Vajramani A., Witham FM, Richards RH (2010) Congenital unilateral absence of sternocleidomastoid and trapezius muscles: a case report and literature review. *J Pediatr Orthop B* 19: 462-4.
16. Bergin M, Elliott J, Jull G (2011) Absence of the inferior portion of the trapezius muscle in three family members. *Manual Therapy* 16: 629-35.
17. Newman CJ, Jacquemont S, Theumann N, Jeann PY(2011) Familial aplasia of the trapezius muscle: clinical and MRI findings. *Acta Paediatr* 100: 464-6.